

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
30 June 2005 (30.06.2005)

PCT

(10) International Publication Number
WO 2005/058874 A1

(51) International Patent Classification⁷: **C07D 401/14**,
A61K 31/4184, A61P 11/00, 31/12, C07D 401/06, 405/14
// (C07D 401/14, 235:00, 213:00, 211:00)

(21) International Application Number:
PCT/EP2004/053618

(22) International Filing Date:
20 December 2004 (20.12.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
03104806.9 18 December 2003 (18.12.2003) EP
60/566,867 30 April 2004 (30.04.2004) US

(71) Applicant (for all designated States except US): **TI-
BOTEC PHARMACEUTICALS LTD.** [IE/IE]; Little
Island, Co Cork (IE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BONFANTI,
Jean-François** [FR/FR]; 4 bis Route Nationale, F-27430
Andé (FR). **ANDRIES, Koenraad, Jozef, Lodewijk**
[BE/BE]; Oosteneinde 9, B-2340 Beerse (BE). **FORTIN,
Jérôme, Michel, Claude** [FR/FR]; 22, rue des acacias,
F-27460 Igoville (FR). **MULLER, Philippe** [FR/FR];
17 Route d'Herqueville, F-27430 Andé (FR). **DOU-
BLET, Frédéric, Marc, Maurice** [FR/FR]; 1646, Route
de Neufchatel, F-76230 Isneauville (FR). **MEYER,
Christophe** [FR/FR]; 428 rue de Gouy, F-76520 Les
Authieux s/l Port St Ouen (FR). **WILLEBRORDS, Rudy,
Edmond** [BE/BE]; Vaartstraat 70, B-2330 Merksplas
(BE). **GEVERS, Tom, Valerius, Josepha** [BE/BE];
Burgemeester Bossaertlaan 17, B-2350 Vosselaar (BE).
TIMMERMAN, Philip, Maria, Bern [BE/BE];
Vijversstraat 211, B-3550 Hasselt (BE).

(74) Agent: **WANTE, Dirk**; Tibotec-Virco Comm. VA, Gen-
eraal De Wittelaan L 11B 3, B-2800 Mechelen (BE).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO,
SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN,
GQ, GW, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted
a patent (Rule 4.17(ii)) for the following designations AE,
AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ,
CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE,
EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS,
JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA,
MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM,
PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY,
TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA,
ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ,
NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM,
AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT,
BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR,
HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG)

— as to the applicant's entitlement to claim the priority of the
earlier application (Rule 4.17(iii)) for all designations

[Continued on next page]

(54) Title: 5- OR 6-SUBSTITUTED BENZIMIDAZOLE DERIVATIVES AS INHIBITORS OF RESPIRATORY SYNCYTIAL
VIRUS REPLICATION



(57) Abstract: The present invention concerns 5- or 6-substituted-benzimidazole deriva-
tives having inhibitory activity on the replication of the respiratory syncytial virus and
having the formula (I) a prodrug, N-oxide, addition salt, quaternary amine, metal complex
or stereochemically isomeric form thereof wherein Q is Ar², R⁶, pyrrolidiny1 substituted
with R⁶, piperidiny1 substituted with R⁶ or homopiperidiny1 substituted with R⁶, G is a di-
rect bond or optionally substituted C₁₋₁₀alkanediyl; R¹ is Ar¹ or a monocyclic or bicyclic
heterocycle; one of R^{2a} and R^{2b} is cyanoC₁₋₆alkyl, cyanoC₂₋₆alkenyl, Ar³C₁₋₆alkyl, Het-
C₁₋₆alkyl, N(R^{8a}R^{8b})C₁₋₆alkyl, Ar³C₂₋₆alkenyl, Het-C₂₋₆alkenyl, Ar³aminoC₁₋₆alkyl, Het-
aminoC₁₋₆alkyl, Ar³thioC₁₋₆alkyl, Het-thioC₁₋₆alkyl, Ar³sulfonylC₁₋₆alkyl, HetsulfonylC₁₋₆
alkyl, Ar³aminocarbonyl, Het-aminocarbonyl, Ar³(CH₂)_naminocarbonyl, Het-(CH₂)_naminocarbonyl, Ar³carbonylamino, Het-car-
bonylamino, Ar³(CH₂)_ncarbonylamino, Het-(CH₂)_ncarbonylamino, and the other one of R^{2a} and R^{2b} is hydrogen; in case R^{2a} is
hydrogen, then R³ is hydrogen; in case R^{2b} is hydrogen, the R³ is hydrogen or C₁₋₆alkyl. It further concerns their preparation and
compositions comprising them, as well as their use as a medicine.



— of inventorship (Rule 4.17(iv)) for US only

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

5- OR 6-SUBSTITUTED BENZIMIDAZOLE DERIVATIVES AS INHIBITORS OF RESPIRATORY SYNCYTIAL VIRUS REPLICATION

The present invention is concerned with 5- or 6-substituted-benzimidazole derivatives having antiviral activity, in particular, having an inhibitory activity on the replication of the respiratory syncytial virus (RSV). It further concerns the preparation thereof and compositions comprising these compounds.

Human RSV or Respiratory Syncytial Virus is a large RNA virus, member of the family of Paramyxoviridae, subfamily pneumovirinae together with bovine RSV virus. Human RSV is responsible for a spectrum of respiratory tract diseases in people of all ages throughout the world. It is the major cause of lower respiratory tract illness during infancy and childhood. Over half of all infants encounter RSV in their first year of life, and almost all within their first two years. The infection in young children can cause lung damage that persists for years and may contribute to chronic lung disease in later life (chronic wheezing, asthma). Older children and adults often suffer from a (bad) common cold upon RSV infection. In old age, susceptibility again increases, and RSV has been implicated in a number of outbreaks of pneumonia in the aged resulting in significant mortality.


Infection with a virus from a given subgroup does not protect against a subsequent infection with an RSV isolate from the same subgroup in the following winter season. Re-infection with RSV is thus common, despite the existence of only two subtypes, A and B.

Today only three drugs have been approved for use against RSV infection. A first one is ribavirin, a nucleoside analogue, provides an aerosol treatment for serious RSV infection in hospitalized children. The aerosol route of administration, the toxicity (risk of teratogenicity), the cost and the highly variable efficacy limit its use. The other two drugs, RespiGam® and palivizumab, polyclonal and monoclonal antibody immunostimulants, are intended to be used in a preventive way.




Other attempts to develop a safe and effective RSV vaccine have all met with failure thus far. Inactivated vaccines failed to protect against disease, and in fact in some cases enhanced disease during subsequent infection. Live attenuated vaccines have been tried with limited success. Clearly there is a need for an efficacious non-toxic and easy to administer drug against RSV replication.

Previously, benzimidazoles and imidazopyridines as inhibitors of RSV replication have been described in WO 01/00611, WO 01/00612 and WO 01/00615.


- Several series of benzimidazolyl and imidazopyridinyl piperidines have been described in patents, patent applications and publications of Janssen Pharmaceutica N.V. as compounds possessing antihistaminic properties. See for example EP-A-5 318, EP-A-99 139, EP-A-145 037, WO-92/01687, Janssens F. et al. in Journal of Medicinal Chemistry, Am. Chem. Soc., Vol. 28, no. 12, pp. 1934-1943 (1985).
- 10 The present invention concerns inhibitors of RSV replication, which can be represented by formula (I)



- their prodrugs, *N*-oxides, addition salts, quaternary amines, metal complexes and stereochemically isomeric forms wherein
- 15 Q is Ar², R^{6a}, pyrrolidinyl substituted with R⁶, piperidinyl substituted with R⁶ or homopiperidinyl substituted with R⁶;
- G is a direct bond or C₁₋₁₀alkanediyl optionally substituted with one or more substituents individually selected from the group consisting of hydroxy, C₁₋₆alkyloxy, Ar¹C₁₋₆alkyloxy, C₁₋₆alkylthio, Ar¹C₁₋₆alkylthio,
- 20 HO(-CH₂-CH₂-O)_n, C₁₋₆alkyloxy(-CH₂-CH₂-O)_n and Ar¹C₁₋₆alkyloxy(-CH₂-CH₂-O)_n;
- R¹ is Ar¹ or a monocyclic or bicyclic heterocycle being selected from piperidinyl, piperazinyl, pyridyl, pyrazinyl, pyridazinyl, pyrimidinyl, furanyl, tetrahydrofuryl, thienyl, pyrrolyl, thiazolyl, oxazolyl, imidazolyl, isothiazolyl, pyrazolyl,
- 25 isoxazolyl, oxadiazolyl, quinolinyl, quinoxalinyl, benzofuranyl, benzothienyl, benzimidazolyl, benzoxazolyl, benzthiazolyl, pyridopyridyl, naphthiridinyl, 1*H*-imidazo[4,5-*b*]pyridinyl, 3*H*-imidazo[4,5-*b*]pyridinyl, imidazo[1,2-*a*]pyridinyl, 2,3-dihydro-1,4-dioxino[2,3-*b*]pyridyl or a radical of formula



-3-



- wherein each of said monocyclic or bicyclic heterocycles may optionally be substituted
- 5 with 1 or where possible more, such as 2, 3, 4 or 5, substituents individually selected from the group of substituents consisting of halo, hydroxy, amino, cyano, carboxyl, C₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkylthio, C₁₋₆alkyloxyC₁₋₆alkyl, Ar¹, Ar¹C₁₋₆alkyl, Ar¹C₁₋₆alkyloxy, hydroxyC₁₋₆alkyl, mono- or di(C₁₋₆alkyl)amino, mono- or di(C₁₋₆alkyl)aminoC₁₋₆alkyl, polyhaloC₁₋₆alkyl, C₁₋₆alkylcarbonylamino,
- 10 C₁₋₆alkyl-SO₂-NR^{4a}-, Ar¹-SO₂-NR^{4a}-, C₁₋₆alkyloxycarbonyl, -C(=O)-NR^{4a}R^{4b}, HO(-CH₂-CH₂-O)_n-, halo(-CH₂-CH₂-O)_n-, C₁₋₆alkyloxy(-CH₂-CH₂-O)_n-, Ar¹C₁₋₆alkyloxy(-CH₂-CH₂-O)_n- and mono- and di(C₁₋₆alkyl)amino(-CH₂-CH₂-O)_n;
- one of R^{2a} and R^{2b} is cyanoC₁₋₆alkyl, cyanoC₂₋₆alkenyl, Ar³C₁₋₆alkyl, (Ar³)(OH)C₁₋₆alkyl, Het-C₁₋₆alkyl, N(R^{8a}R^{8b})C₁₋₆alkyl, Ar³C₂₋₆alkenyl,
- 15 Het-C₂₋₆alkenyl, Ar³aminoC₁₋₆alkyl, Het-aminoC₁₋₆alkyl, Het-C₁₋₆alkylamino-C₁₋₆alkyl, Ar³thioC₁₋₆alkyl, Het-thioC₁₋₆alkyl, Ar³sulfonylC₁₋₆alkyl, Het-sulfonyl-C₁₋₆alkyl, Ar³aminocarbonyl, Het-aminocarbonyl, Ar³(CH₂)_naminocarbonyl, Het-(CH₂)_naminocarbonyl, Ar³carbonylamino, Het-carbonylamino, Ar³(CH₂)_ncarbonylamino, Het-(CH₂)_ncarbonylamino or Ar³(CH₂)_namino; and the
- 20 other one of R^{2a} and R^{2b} is hydrogen;
- in case R^{2a} is hydrogen, then R³ is hydrogen;
- in case R^{2b} is hydrogen, then R³ is hydrogen or C₁₋₆alkyl;
- R^{4a} and R^{4b} can be the same or can be different relative to one another, and are each independently hydrogen or C₁₋₆alkyl; or
- 25 R^{4a} and R^{4b} taken together may form a bivalent radical of formula -(CH₂)_s- wherein s is 4 or 5;
- R⁵ is hydrogen or C₁₋₆alkyl;
- R⁶ is hydrogen or C₁₋₆alkyl optionally substituted with one or more substituents each independently selected from the group consisting of trifluoromethyl, NR^{7a}R^{7b}, C₃₋₇cycloalkyl, Ar², hydroxy, C₁₋₄alkoxy, C₁₋₄alkylthio, Ar²-oxy-, Ar²-thio-, Ar²(CH₂)_noxy, Ar²(CH₂)_nthio, hydroxycarbonyl, aminocarbonyl, C₁₋₄alkyl-
- 30

- carbonyl, Ar²carbonyl, C₁₋₄alkoxycarbonyl, Ar²(CH₂)_ncarbonyl, aminocarbonyloxy, C₁₋₄alkylcarbonyloxy, Ar²carbonyloxy, Ar²(CH₂)_ncarbonyloxy, C₁₋₄alkoxy-carbonyl(CH₂)_noxy, mono- or di(C₁₋₄alkyl)aminocarbonyl, mono- or di(C₁₋₄alkyl)-aminocarbonyloxy, aminosulfonyl, mono- or di(C₁₋₄alkyl)aminosulfonyl or a
- 5 heterocycle selected from the group consisting of pyrrolidinyl, pyrrolyl, dihydropyrrolyl, imidazolyl, triazolyl, piperidinyl, homopiperidinyl, piperazinyl, dioxolanyl, dioxanyl, pyridyl and tetrahydropyridyl, wherein each of said heterocycle may optionally be substituted with one or two radicals selected from oxo and C₁₋₆alkyl;
- 10 R^{6a} is C₁₋₆alkyl substituted with one or more substituents each independently selected from the group consisting of trifluoromethyl, NR^{7a}R^{7b}, C₃₋₇cycloalkyl, Ar², hydroxy, C₁₋₄alkoxy, C₁₋₄alkylthio, Ar²-oxy-, Ar²-thio-, Ar²(CH₂)_noxy, Ar²(CH₂)_nthio, hydroxycarbonyl, aminocarbonyl, C₁₋₄alkylcarbonyl, Ar²carbonyl, C₁₋₄alkoxycarbonyl, Ar²(CH₂)_ncarbonyl, aminocarbonyloxy, C₁₋₄alkylcarbonyloxy, Ar²carbonyloxy, Ar²(CH₂)_ncarbonyloxy, C₁₋₄alkoxycarbonyl(CH₂)_noxy,
- 15 mono- or di(C₁₋₄alkyl)aminocarbonyl, mono- or di(C₁₋₄alkyl)aminocarbonyloxy, aminosulfonyl, mono- or di(C₁₋₄alkyl)aminosulfonyl or a heterocycle selected from the group consisting of pyrrolidinyl, pyrrolyl, dihydropyrrolyl, imidazolyl, triazolyl, piperidinyl, homopiperidinyl, piperazinyl, dioxolanyl, dioxanyl, pyridyl and tetrahydropyridyl, wherein each of said heterocycle may optionally be
- 20 substituted with one or two radicals selected oxo and C₁₋₆alkyl;
- R^{7a} is hydrogen, C₁₋₆alkyl, formyl or C₁₋₆alkylcarbonyl;
- R^{7b} is hydrogen, C₁₋₆alkyl, formyl or C₁₋₆alkylcarbonyl;
- R^{8a} is Ar³, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, C₁₋₆alkoxyC₁₋₆alkyl, cyanoC₁₋₆alkyl,
- 25 aminoC₁₋₆alkyl, mono-or di(C₁₋₆alkyl)aminoC₁₋₆alkyl, Ar³C₁₋₆alkyl, Het-C₁₋₆alkyl, aminocarbonyl-C₁₋₆-alkyl, carboxyl-C₁₋₆-alkyl;
- R^{8b} is Ar³, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, C₁₋₆alkoxyC₁₋₆alkyl, cyanoC₁₋₆alkyl, aminoC₁₋₆alkyl, mono-or di(C₁₋₆alkyl)aminoC₁₋₆alkyl, Ar³C₁₋₆alkyl, Het-C₁₋₆alkyl;
- each n independently is 1, 2, 3 or 4;
- 30 each m independently is 1 or 2;
- each p independently is 1 or 2;
- Ar¹ is phenyl or phenyl substituted with 1 or more, such as 2, 3 or 4, substituents selected from halo, hydroxy, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, polyhaloC₁₋₆alkyl, and C₁₋₆alkyloxy;
- 35 Ar² is phenyl or phenyl substituted with 1 or more, such as 2, 3 or 4, substituents selected from the group consisting of halo, hydroxy, amino, cyano, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, polyhaloC₁₋₆alkyl, aminoC₁₋₆alkyl, C₁₋₆alkyloxy, aminosulfonyl, aminocarbonyl, hydroxycarbonyl, C₁₋₄alkylcarbonyl, mono- or

di(C₁₋₄alkyl)amino, mono- or di(C₁₋₄alkyl)aminocarbonyl, mono- or di(C₁₋₄alkyl)aminosulfonyl, mono- or di(C₁₋₄alkyl)aminoC₁₋₆alkyl and C₁₋₄alkoxycarbonyl;

- Ar³ is phenyl, naphthalenyl, 1,2,3,4-tetrahydro-naphthalenyl or indanyl, wherein said
 5 phenyl, naphthyl, 1,2,3,4-tetrahydro-naphthalenyl or indanyl may optionally and each individually be substituted with one or more, such as 2, 3 or 4, substituents selected from the group consisting of halo, hydroxy, mercapto, amino, cyano, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, Ar¹, hydroxyC₁₋₆alkyl, polyhaloC₁₋₆alkyl, aminoC₁₋₆alkyl, cyanoC₁₋₆alkyl, aminocarbonyl, C₁₋₆alkyloxy, C₁₋₆alkylthio,
 10 Ar¹-oxy, Ar¹-thio, Ar¹-amino, aminosulfonyl, aminocarbonyl-C₁₋₆alkyl, hydroxyl-carbonyl-C₁₋₆alkyl, hydroxycarbonyl, C₁₋₄alkylcarbonyl, mono- or di(C₁₋₄alkyl)amino, mono- or di(C₁₋₄alkyl)aminocarbonyl, mono- or di(C₁₋₄alkyl)aminosulfonyl, mono- or di(C₁₋₄alkyl)aminoC₁₋₆alkyl, C₁₋₄alkylcarbonylamino and C₁₋₄alkoxycarbonyl;
 15 Het is a heterocycle being selected from tetrahydrofuranyl, tetrahydrothienyl, dioxanyl, dioxolanyl, pyrrolidinyl, pyrrolidinonyl, furanyl, thienyl, pyrrolyl, thiazolyl, oxazolyl, imidazolyl, isothiazolyl, pyrazolyl, isoxazolyl, oxadiazolyl, thiadiazolyl, piperidinyl, homopiperidinyl, piperazinyl, morpholinyl, pyridyl, pyrazinyl, pyridazinyl, pyrimidinyl, tetrahydroquinolinyl, quinolinyl, isoquinolinyl,
 20 benzodioxanyl, benzodioxolyl, indolinyl, indolyl, each of said heterocycle may optionally be substituted with oxo, amino, Ar¹, C₁₋₄alkyl, aminoC₁₋₄alkyl, hydroxyC₁₋₆alkyl, Ar¹C₁₋₄alkyl, mono- or di(C₁₋₆alkyl)aminoC₁₋₆alkyl, mono- or di(C₁₋₆alkyl)amino, or with two C₁₋₄alkyl radicals.

- 25 The invention also relates to the use of a compound of formula (I), or a prodrug, *N*-oxide, addition salt, quaternary amine, metal complex and stereochemically isomeric form thereof, for the manufacture of a medicament for inhibiting RSV replication. Or the invention relates to a method of inhibiting RSV replication in a warm-blooded animal said method comprising the administration of an effective amount of a
 30 compound of formula (I), or a prodrug, *N*-oxide, addition salt, quaternary amine, metal complex and stereochemically isomeric form thereof.

- In a further aspect, this invention relates to novel compounds of formula (I) as well as methods for preparing these compounds.
 35

The term prodrug as used throughout this specification and claims means the pharmacologically acceptable derivatives, e.g. esters and amides, such that the resulting biotransformation product of the derivative is the active drug as defined in the

compounds of formula (I). The reference by Goodman and Gilman (The Pharmacological Basis of Therapeutics, 8th ed., McGraw-Hill, Int. Ed. 1992, "Biotransformation of Drugs", p. 13-15) describing prodrugs generally, is hereby incorporated. Prodrugs are characterized by a good aqueous solubility and
5 bioavailability, and are readily metabolized into the active inhibitors *in vivo*.

The terms 'C₁₋₁₀alkanediyl optionally substituted with one or more substituents' and 'C₁₋₆alkyl optionally substituted with one or more substituents' such as used in the definition of G and respectively R⁶ or R^{6a} are meant to comprise C₁₋₁₀alkanediyl or
10 C₁₋₆alkyl radicals having two or more substituents, for example two, three, four, five or six substituents, in particular two or three substituents, further in particular two substituents. The upper limit of the number of substituents is determined by the number of hydrogen atoms that can be replaced as well as by the general properties of the substituents such as their bulkiness, these properties allowing the skilled person to
15 determine said upper limit.

As used in the foregoing and hereinafter, 'polyhaloC₁₋₆alkyl' as a group or part of a group, e.g. in polyhaloC₁₋₆alkyloxy, is defined as mono- or polyhalo substituted C₁₋₆alkyl, in particular C₁₋₆alkyl substituted with up to one, two, three, four, five, six, or
20 more halo atoms, such as methyl or ethyl with one or more fluoro atoms, for example, difluoromethyl, trifluoromethyl, trifluoroethyl. Also included are perfluoro C₁₋₆alkyl groups, which are C₁₋₆alkyl groups whereon all hydrogen atoms are replaced by fluoro atoms, e.g. pentafluoroethyl. In case more than one halogen atom is attached to an alkyl group within the definition of polyhaloC₁₋₆alkyl, the halogen atoms may be the same or
25 different.

Each of the monocyclic or bicyclic heterocycles in the definition of R¹ may optionally be substituted with 1 or where possible more substituents, such as 2, 3, 4 or 5, substituents. In particular, said heterocycles may optionally be substituted with up to 4,
30 up to 3, up to 2 substituents, or up to 1 substituent.

Each Ar¹ or Ar² may be unsubstituted phenyl or phenyl substituted with 1 or more substituents, such as 5 or 4 substituents or, which is preferred, up to 3 substituents, or up to two substituents, or with one substituent. Ar³ is phenyl, naphthalenyl, 1,2,3,4-
35 tetrahydro-naphthalenyl or indanyl, which each may optionally with one or more substituents, such as 5 or 4 substituents or, which is preferred, up to 3 substituents, or up to two substituents, or with one substituent.

A hydroxyC₁₋₆alkyl group when substituted on an oxygen atom or a nitrogen atom preferably is a hydroxyC₂₋₆alkyl group wherein the hydroxy group and the oxygen or nitrogen are separated by at least two carbon atoms.

- 5 R⁶ or R^{6a} can be C₁₋₆alkyl substituted with one or more substituents selected from NR^{7a}R^{7b}, hydroxy, C₁₋₄alkoxy, C₁₋₄alkylthio, Ar²-oxy-, Ar²-thio-, Ar²(CH₂)_noxy, Ar²(CH₂)_nthio, aminocarbonyloxy, C₁₋₄alkylcarbonyloxy, Ar²carbonyloxy, Ar²(CH₂)_ncarbonyloxy, C₁₋₄alkoxycarbonyl(CH₂)_noxy, mono- and di(C₁₋₄alkyl)amino-carbonyloxy. In that instance C₁₋₆alkyl preferably has at least two carbon atoms (i.e.
- 10 C₂₋₆alkyl) and the said substituents are not substituted on the carbon atom linked to the nitrogen bearing Q.

- As used herein C₁₋₄alkyl as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 1 to 4 carbon atoms such as methyl, ethyl,
- 15 propyl, 1-methylethyl, butyl and the like; C₂₋₄alkyl as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 2 to 4 carbon atoms such as ethyl, propyl, 1-methylethyl, butyl and the like; C₁₋₆alkyl as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 1 to 6 carbon atoms such as the groups defined for C₁₋₄alkyl and pentyl,
- 20 hexyl, 2-methylbutyl and the like.

- As used herein C₂₋₆alkenyl as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having at least one double bond, and preferably having one double bond, and further having from 2 to 6 carbon atoms such as ethenyl,
- 25 propenyl, buten-1-yl, buten-2-yl, penten-1-yl, penten-2-yl, hexen-1-yl, hexen-2-yl, hexen-3-yl, 2-methylbuten-1-yl and the like.

- C₃₋₇cycloalkyl is generic to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl.
- 30

- C₂₋₅alkanediyl defines bivalent straight and branched chain saturated hydrocarbon radicals having from 2 to 5 carbon atoms such as, for example, 1,2-ethanediyl, 1,3-propanediyl, 1,4-butanediyl, 1,2-propanediyl, 2,3-butanediyl, 1,5-pentanediyl and the like, C₁₋₄alkanediyl defines bivalent straight and branched chain saturated
- 35 hydrocarbon radicals having from 1 to 4 carbon atoms such as, for example, methylene, 1,2-ethanediyl, 1,3-propanediyl, 1,4-butanediyl and the like; C₁₋₆alkanediyl is meant to include C₁₋₄alkanediyl and the higher homologues thereof having from 5 to 6 carbon atoms such as, for example, 1,5-pentanediyl, 1,6-hexanediyl and the like; C₁₋₁₀alkanediyl is meant to include C₁₋₆alkanediyl and the higher homologues thereof

having from 7 to 10 carbon atoms such as, for example, 1,7-heptanediyl, 1,8-octanediyl, 1,9-nonanediyl, 1,10-decanediyl and the like.

As used herein before, the term (=O) forms a carbonyl moiety when attached to a carbon atom, a sulfoxide moiety when attached to a sulfur atom and a sulfonyl moiety when two of said terms are attached to a sulfur atom. The term (=N-OH) forms a hydroxyimine moiety when attached to a carbon atom.

The term halo is generic to fluoro, chloro, bromo and iodo.

It should be noted that the radical positions on any molecular moiety used in the definitions may be anywhere on such moiety as long as it is chemically stable.

Radicals used in the definitions of the variables include all possible isomers unless otherwise indicated. For instance pyridyl includes 2-pyridyl, 3-pyridyl and 4-pyridyl; pentyl includes 1-pentyl, 2-pentyl and 3-pentyl.

When any variable occurs more than one time in any constituent, each definition is independent.

Whenever used hereinafter, the term "compounds of formula (I)", or "the present compounds" or similar term is meant to include the compounds of general formula (I), their prodrugs, *N*-oxides, addition salts, quaternary amines, metal complexes and stereochemically isomeric forms. An interesting subgroup of the compounds of formula (I) or any subgroup thereof are the *N*-oxides, salts and all the stereoisomeric forms of the compounds of formula (I).

It will be appreciated that some of the compounds of formula (I) may contain one or more centers of chirality and exist as stereochemically isomeric forms.

The term "stereochemically isomeric forms" as used hereinbefore defines all the possible compounds made up of the same atoms bonded by the same sequence of bonds but having different three-dimensional structures which are not interchangeable, which the compounds of formula (I) may possess.

Unless otherwise mentioned or indicated, the chemical designation of a compound encompasses the mixture of all possible stereochemically isomeric forms which said compound may possess. Said mixture may contain all diastereomers and/or enantiomers of the basic molecular structure of said compound. All stereochemically isomeric

forms of the compounds of the present invention both in pure form or in admixture with each other are intended to be embraced within the scope of the present invention.

Pure stereoisomeric forms of the compounds and intermediates as mentioned herein are defined as isomers substantially free of other enantiomeric or diastereomeric forms of the same basic molecular structure of said compounds or intermediates. In particular, the term 'stereoisomerically pure' concerns compounds or intermediates having a stereoisomeric excess of at least 80% (i.e. minimum 90% of one isomer and maximum 10% of the other possible isomers) up to a stereoisomeric excess of 100% (i.e. 100% of one isomer and none of the other), more in particular, compounds or intermediates having a stereoisomeric excess of 90% up to 100%, even more in particular having a stereoisomeric excess of 94% up to 100% and most in particular having a stereoisomeric excess of 97% up to 100%. The terms 'enantiomerically pure' and 'diastereomerically pure' should be understood in a similar way, but then having regard to the enantiomeric excess, respectively the diastereomeric excess of the mixture in question.

Pure stereoisomeric forms of the compounds and intermediates of this invention may be obtained by the application of art-known procedures. For instance, enantiomers may be separated from each other by the selective crystallization of their diastereomeric salts with optically active acids or bases. Examples thereof are tartaric acid, dibenzoyl-tartaric acid, ditoluoyltartaric acid and camphosulfonic acid. Alternatively, enantiomers may be separated by chromatographic techniques using chiral stationary phases. Said pure stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting materials, provided that the reaction occurs stereospecifically. Preferably, if a specific stereoisomer is desired, said compound will be synthesized by stereospecific methods of preparation. These methods will advantageously employ enantiomerically pure starting materials.

The diastereomeric racemates of formula (I) can be obtained separately by conventional methods. Appropriate physical separation methods that may advantageously be employed are, for example, selective crystallization and chromatography, e.g. column chromatography.

For some of the compounds of formula (I), their prodrugs, N-oxides, salts, solvates, quaternary amines, or metal complexes and the intermediates used in the preparation thereof, the absolute stereochemical configuration was not experimentally determined.

A person skilled in the art is able to determine the absolute configuration of such compounds using art-known methods such as, for example, X-ray diffraction.

5 The present invention is also intended to include all isotopes of atoms occurring on the present compounds. Isotopes include those atoms having the same atomic number but different mass numbers. By way of general example and without limitation, isotopes of hydrogen include tritium and deuterium. Isotopes of carbon include C-13 and C-14.

10 For therapeutic use, salts of the compounds of formula (I) are those wherein the counterion is pharmaceutically acceptable. However, salts of acids and bases which are non-pharmaceutically acceptable may also find use, for example, in the preparation or purification of a pharmaceutically acceptable compound. All salts, whether pharmaceutically acceptable or not are included within the ambit of the present invention.

15 The pharmaceutically acceptable acid and base addition salts as mentioned hereinabove are meant to comprise the therapeutically active non-toxic acid and base addition salt forms which the compounds of formula (I) are able to form. The pharmaceutically acceptable acid addition salts can conveniently be obtained by treating the base form with such appropriate acid. Appropriate acids comprise, for example, inorganic acids
20 such as hydrohalic acids, e.g. hydrochloric or hydrobromic acid, sulfuric, nitric, phosphoric and the like acids; or organic acids such as, for example, acetic, propanoic, hydroxyacetic, lactic, pyruvic, oxalic (i.e. ethanedioic), malonic, succinic (i.e. butanedioic acid), maleic, fumaric, malic (i.e. hydroxybutanedioic acid), tartaric, citric, methanesulfonic, ethanesulfonic, benzenesulfonic, *p*-toluenesulfonic, cyclamic,
25 salicylic, *p*-aminosalicylic, pamoic and the like acids.

Conversely said salt forms can be converted by treatment with an appropriate base into the free base form.

30 The compounds of formula (I) containing an acidic proton may also be converted into their non-toxic metal or amine addition salt forms by treatment with appropriate organic and inorganic bases. Appropriate base salt forms comprise, for example, the ammonium salts, the alkali and earth alkaline metal salts, e.g. the lithium, sodium, potassium, magnesium, calcium salts and the like, salts with organic bases, e.g. the
35 benzathine, *N*-methyl-D-glucamine, hydrabamine salts, and salts with amino acids such as, for example, arginine, lysine and the like.

The term addition salt as used hereinabove also comprises the solvates which the compounds of formula (I) as well as the salts thereof, are able to form. Such solvates are for example hydrates, alcoholates and the like.

- 5 The term "quaternary amine" as used hereinbefore defines the quaternary ammonium salts which the compounds of formula (I) are able to form by reaction between a basic nitrogen of a compound of formula (I) and an appropriate quaternizing agent, such as, for example, an optionally substituted alkylhalide, arylhalide or arylalkylhalide, e.g. methyl iodide or benzyl iodide. Other reactants with good leaving groups may also be used, such as alkyl trifluoromethanesulfonates, alkyl methanesulfonates, and alkyl p-toluenesulfonates. A quaternary amine has a positively charged nitrogen. Pharmaceutically acceptable counterions include chloro, bromo, iodo, trifluoroacetate and acetate. The counterion of choice can be introduced using ion exchange resins.
- 10
- 15 The *N*-oxide forms of the present compounds are meant to comprise the compounds of formula (I) wherein one or several nitrogen atoms are oxidized to the so-called *N*-oxide.

It will be appreciated that the compounds of formula (I) may have metal binding, chelating, complex forming properties and therefore may exist as metal complexes or metal chelates. Such metalated derivatives of the compounds of formula (I) are intended to be included within the scope of the present invention.

20


Some of the compounds of formula (I) may also exist in their tautomeric form. Such forms although not explicitly indicated in the above formula are intended to be included within the scope of the present invention.

25

Any subgroup of compounds of formula (I) specified herein is meant to also comprise the prodrugs, *N*-oxides, addition salts, quaternary amines, metal complexes and stereochemically isomeric forms of this subgroup of compounds of formula (I).

30

One embodiment of the present invention concerns compounds of formula (I-a):




wherein Q, R⁵, G, R¹ and R^{2b} are as specified above in the definitions of the compounds of formula (I) or as in any of the subgroups of compounds specified herein.

35


Another embodiment of the present invention concerns compounds of formula (I-b):

-12-



wherein Q, R⁵, G, R¹, R^{2a} and R³ are as specified above in the definitions of the compounds of formula (I) or as in any of the subgroups of compounds specified herein.

- 5 One particular embodiment of the present invention concerns compounds of formula (I-a-1):



wherein Q, R⁵, G and R¹ are as specified in the definitions of the compounds of formula (I) or any of the subgroups of compounds of formula (I) specified herein; and


- 10 Alk is C₁₋₆alkanediyl;

R^{8c} has the same meanings of R^{8a}, and also may be hydrogen;

R⁹, R¹⁰, R¹¹ independently from one another have the same meanings as the substituents on Ar³ as specified in the definitions of the compounds of formula (I) or of any of the subgroups thereof.

15

Another particular embodiment of the present invention concerns compounds of formula (I-b-1):



wherein Q, R⁵, G, R¹ and R³ are as specified in the definitions of the compounds of formula (I) or any of the subgroups of compounds of formula (I) specified herein; and Alk is C₁₋₆alkanediyl;


20

R^{8c} has the same meanings of R^{8a}, and also may be hydrogen;

R⁹, R¹⁰, R¹¹ independently from one another have the same meanings as the substituents on Ar³ as specified in the definitions of the compounds of formula (I) or of any of the subgroups thereof.

25


Another embodiment of the present invention concerns compounds of formula (I-c):



wherein t , G , R^1 , R^{2a} , R^{2b} , R^3 , R^5 and R^6 are as specified above in the definition of the compounds of formula (I), or as in any of the subgroups of compounds specified

5 herein.


Another embodiment of the present invention concerns compounds of formula (I-d):



wherein t , R^5 , R^6 , G , R^1 and R^{2b} are as specified above or as in any of the subgroups of compounds specified herein.

10


Another embodiment of the present invention concerns compounds of formula (I-e):





wherein t , R^5 , R^6 , G , R^1 , R^{2a} and R^3 are as specified above or as in any of the subgroups of compounds specified herein.

15

Still further embodiments comprise compounds of formula (I-c), (I-d) or (I-e) wherein t is 2, i.e. compounds of formulae




-14-




wherein Q, t, R⁵, G, R¹, R^{2a}, R^{2b}, R³ are as specified above or as in any of the subgroups
 5 of compounds specified herein.

Another embodiment of the present invention concerns compounds of formula (I-d-2):



wherein R⁵, R⁶, G and R¹ are as specified above or as in any of the subgroups of
 10 compounds specified herein; and
 t is 1, 2 or 3; preferably t is 2;
 Alk is C₁₋₆alkanediyl;
 R^{8c} has the same meanings of R^{8a}, and also may be hydrogen;
 R⁹, R¹⁰, R¹¹ independently from one another have the same meanings as the
 15 substituents on Ar³ as specified in the definitions of the compounds of formula (I)
 or of any of the subgroups thereof.

Another embodiment of the present invention concerns compounds of formula (I-e):



20 wherein R⁵, R⁶, G, R¹ and R³ are as specified above or as in any of the subgroups of

compounds specified herein; and

t is 1, 2 or 3; preferably t is 2;

R^{8c} has the same meanings of R^{8a}, and also may be hydrogen;

R⁹, R¹⁰, R¹¹ independently from one another have the same meanings as the

- 5 substituents on Ar³ as specified in the definitions of the compounds of formula (I) or of any of the subgroups thereof.

Further preferred subgroups are those wherein Alk is ethylene or methylene, more preferably wherein Alk is methylene.

10

In (I-a-1), (I-b-1), (I-d-2) or (I-e-2) R^{8c} preferably is hydrogen, hydroxyC₁₋₆alkyl, aminocarbonyl-C₁₋₆alkyl.

In (I-a-1), (I-b-1), (I-d-2) or (I-e-2):

- 15 (a) R⁹, R¹⁰, R¹¹ preferably and independently from one another are hydrogen, halo, hydroxy, mercapto, amino, cyano, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, Ar¹, hydroxyC₁₋₆alkyl, CF₃, aminoC₁₋₆alkyl, cyanoC₁₋₆alkyl, aminocarbonyl, C₁₋₆alkyloxy, C₁₋₆alkylthio, Ar¹-oxy, Ar¹-thio, Ar¹-amino, aminosulfonyl, aminocarbonyl-C₁₋₆alkyl, hydroxycarbonyl-C₁₋₆alkyl, hydroxycarbonyl, C₁₋₄alkylcarbonyl, C₁₋₄alkylcarbonylamino or C₁₋₄alkoxycarbonyl; or
- 20 (b) R⁹, R¹⁰, R¹¹ more preferably and independently from one another are hydrogen, halo, hydroxy, mercapto, cyano, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, Ar¹, hydroxy-C₁₋₆alkyl, CF₃, aminoC₁₋₆alkyl, cyanoC₁₋₆alkyl, aminocarbonyl, C₁₋₆alkyloxy, C₁₋₆alkylthio, Ar¹-oxy, Ar¹-thio, Ar¹-amino, aminosulfonyl, aminocarbonyl-C₁₋₆alkyl, hydroxycarbonyl-C₁₋₆alkyl, C₁₋₄alkylcarbonyl, C₁₋₄alkylcarbonylamino or C₁₋₄alkoxycarbonyl; or
- 25 (c) R⁹, R¹⁰, R¹¹ more preferably and independently from one another are halo, C₁₋₆alkyl or hydroxyC₁₋₆alkyl; or
- (d) R⁹, R¹⁰ more preferably are as in (a), (b) or (c) and R¹¹ is hydrogen; or
- 30 (e) R⁹, R¹⁰ more preferably and independently from one another are C₁₋₆alkyl or hydroxy-C₁₋₆alkyl; and R¹¹ is hydrogen; or
- (f) R⁹, R¹⁰ still more preferably are C₁₋₆alkyl and R¹¹ is hydrogen; or
- (g) R⁹ is C₁₋₆alkyl, R¹⁰ is hydroxy-C₁₋₆alkyl and R¹¹ is hydrogen.

- 35 It is to be understood that the above defined subgroups of compounds of formulae (I-a), (I-b), etc. as well as any other subgroup defined herein, are meant to also comprise any prodrugs, N-oxides, addition salts, quaternary amines, metal complexes and stereochemically isomeric forms of such compounds.

Particular subgroups of the compounds of formula (I) are those compounds of formula (I), or any subgroup of compounds of formula (I) specified herein, wherein G is C₁₋₁₀alkanediyl, more in particular wherein G is methylene.

5

Other particular subgroups of the compounds of formula (I) are those compounds of formula (I), or any subgroup of compounds of formula (I) specified herein, wherein

(a) R¹ is other than Ar¹; or wherein

(b) R¹ is Ar¹ or a monocyclic heterocycle, which is as specified in the definitions of the compounds of formula (I) or any of the subgroups thereof.

10

Further particular subgroups of the compounds of formula (I) are those compounds of formula (I), or any subgroup of compounds of formula (I) specified herein, wherein

(c) R¹ is pyridyl optionally substituted with 1 or 2 substituents independently selected from the group consisting of halo, hydroxy, amino, cyano, carboxyl, C₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkylthio, C₁₋₆alkyloxyC₁₋₆alkyl, Ar¹, Ar¹C₁₋₆alkyl, Ar¹C₁₋₆alkyloxy, hydroxyC₁₋₆alkyl, mono- or di(C₁₋₆alkyl)amino, mono- or di(C₁₋₆alkyl)aminoC₁₋₆alkyl, polyhaloC₁₋₆alkyl, C₁₋₆alkylcarbonylamino, C₁₋₆alkyl-SO₂-NR^{4a}-, Ar¹-SO₂-NR^{4a}-, C₁₋₆alkyloxycarbonyl, -C(=O)-NR^{4a}R^{4b}, HO(-CH₂-CH₂-O)_n-, halo(-CH₂-CH₂-O)_n-, C₁₋₆alkyloxy(-CH₂-CH₂-O)_n-, Ar¹C₁₋₆alkyloxy(-CH₂-CH₂-O)_n- and mono- or di(C₁₋₆alkyl)amino(-CH₂-CH₂-O)_n-; or more in particular

15

20

25

30

(d) R¹ is pyridyl substituted with 1 or 2 substituents independently selected from the group consisting of hydroxy, C₁₋₆alkyl, halo, C₁₋₆alkyloxy, Ar¹C₁₋₆alkyloxy and (C₁₋₆alkyloxy)C₁₋₆alkyloxy; preferably wherein

(e) R¹ is pyridyl substituted with 1 or 2 substituents independently selected from the group consisting of hydroxy, C₁₋₆alkyl, halo and C₁₋₆alkyloxy; or wherein

(f) R¹ is pyridyl substituted with 1 or 2 substituents independently selected from the group consisting of hydroxy and C₁₋₆alkyl; more preferably wherein

(g) R¹ is pyridyl substituted with hydroxy and C₁₋₆alkyl; or more preferably wherein


(h) R¹ is pyridyl substituted with hydroxy and methyl; or wherein

(i) R¹ is 3-hydroxy-6-methylpyrid-2-yl.

Further embodiments comprise those compounds of formula (I) or any of the subgroups of compounds of formula (I) wherein

35

(j) R¹ is Ar¹, quinolinyl, benzimidazolyl, a radical of formula



(c-4)

pyrazinyl, or pyridyl; or wherein

- (k) R^1 is Ar^1 , quinolinyl, benzimidazolyl or a radical of formula (c-4) wherein m is 2, pyrazinyl or pyridyl;
- 5 wherein each of the radicals in (j) and (k) may optionally be substituted with the substituents specified in the definition of the compounds of formula (I) and in particular pyridyl may be substituted as specified above in (a) to (i); or more specifically wherein
- 10 (l) R^1 is Ar^1 , quinolinyl, benzimidazolyl or a radical of formula (c-4) wherein m is 2, pyrazinyl or pyridyl, wherein each of these radicals may optionally be substituted with one, two or three radicals selected from the group consisting of halo, hydroxy, C_{1-6} alkyl, C_{1-6} alkyloxy, Ar^1C_{1-6} alkyloxy, $(C_{1-6}$ alkyloxy) C_{1-6} alkyloxy; or more specifically wherein
- 15 (m) R^1 is Ar^1 , quinolinyl, benzimidazolyl or a radical of formula (c-4) wherein m is 2, pyrazinyl or pyridyl, wherein each of these radicals may optionally be substituted with one, two or three radicals selected from the group consisting of halo, hydroxy, C_{1-6} alkyl, C_{1-6} alkyloxy, benzyloxy; or more specifically wherein
- 20 (n) R^1 is phenyl optionally substituted with one, two or three radicals selected from the group consisting of halo, hydroxy, C_{1-6} alkyl, C_{1-6} alkyloxy; quinolinyl; a radical (c-4) wherein m is 2, optionally substituted with up to two radicals selected from C_{1-6} alkyl; benzimidazolyl optionally substituted with C_{1-6} alkyl; pyridyl optionally substituted with one or two radicals selected from hydroxy, halo, C_{1-6} alkyl, benzyloxy and C_{1-6} alkyloxy, pyrazinyl optionally substituted with up to three
- 25 radicals selected from C_{1-6} alkyl; or pyridyl substituted or optionally substituted as specified above in (a) – (i); or wherein
- (o) R^1 is phenyl optionally substituted with one or two radicals selected from the group consisting of halo, hydroxy, C_{1-6} alkyl, C_{1-6} alkyloxy;
- (p) R^1 is quinolinyl;
- 30 (q) R^1 is a radical (c-4) wherein m is 2, optionally substituted with up to two radicals selected from C_{1-6} alkyl;
- (r) R^1 is benzimidazolyl optionally substituted with C_{1-6} alkyl; pyridyl optionally substituted with one or two radicals selected from hydroxy, halo, C_{1-6} alkyl, benzyloxy and C_{1-6} alkyloxy,

- (s) R^1 is pyrazinyl optionally substituted with up to three radicals selected from C_{1-6} alkyl.

Preferred subgroups of compounds of formula (I) or any of the subgroups of compounds of formula (I) are those wherein G is a direct bond or methylene and R^1 is as specified above in (a) – (s). Further preferred are the compounds of formula (I) or any of the subgroups specified herein wherein G is a direct bond and R^1 is a radical (c-4), in particular wherein m is 2, optionally substituted with up to two radicals selected from C_{1-6} alkyl. Further preferred are the compounds of formula (I) or any of the subgroups specified herein wherein G is methylene and R^1 is as specified above in (a) – (s), but is other than a radical (c-4).

- A particular embodiment of the present invention concerns compounds of formula (I) or of any of the subgroups of compounds of formula (I) specified herein wherein
- 15 (a) one of R^{2a} and R^{3a} is selected from cyano C_{1-6} alkyl, cyano C_{2-6} alkenyl, Ar^3C_{1-6} alkyl, $(Ar^3)(OH)C_{1-6}$ alkyl, Het- C_{1-6} alkyl, $N(R^{8a}R^{8b})C_{1-6}$ alkyl, Ar^3C_{2-6} alkenyl, Het- C_{2-6} alkenyl, Ar^3 amino C_{1-6} alkyl, Het-amino C_{1-6} alkyl, Het- C_{1-6} alkylamino- C_{1-6} alkyl, Ar^3 thio C_{1-6} alkyl, Het-thio C_{1-6} alkyl, Ar^3 sulfonyl C_{1-6} alkyl, Het-sulfonyl- C_{1-6} alkyl, Ar^3 aminocarbonyl, Het-aminocarbonyl, $Ar^3(CH_2)_n$ aminocarbonyl, Het- $(CH_2)_n$ aminocarbonyl, Ar^3 carbonylamino, $Ar^3(CH_2)_n$ amino; and the other one of R^{2a} and R^{2b} is hydrogen; or
 - 20 (b) one of R^{2a} and R^{3a} is selected from cyano C_{1-6} alkyl, cyano C_{2-6} alkenyl, Ar^3C_{1-6} alkyl, $(Ar^3)(OH)C_{1-6}$ alkyl, Het- C_{1-6} alkyl, $N(R^{8a}R^{8b})C_{1-6}$ alkyl, Ar^3C_{2-6} alkenyl, Ar^3 amino C_{1-6} alkyl, Het-amino C_{1-6} alkyl, Het- C_{1-6} alkylamino C_{1-6} alkyl, Ar^3 thio- C_{1-6} alkyl, Ar^3 aminocarbonyl, Het-aminocarbonyl, $Ar^3(CH_2)_n$ aminocarbonyl, Het- $(CH_2)_n$ aminocarbonyl, Ar^3 carbonylamino, $Ar^3(CH_2)_n$ amino; and the other one of R^{2a} and R^{2b} is hydrogen; or
 - 25 (c) one of R^{2a} and R^{3a} is selected from cyano C_{1-6} alkyl, Ar^3C_{1-6} alkyl, Het- C_{1-6} alkyl, $N(R^{8a}R^{8b})C_{1-6}$ alkyl, Ar^3C_{2-6} alkenyl, Ar^3 amino C_{1-6} alkyl, Het-amino C_{1-6} alkyl, Het- C_{1-6} alkylamino C_{1-6} alkyl, Ar^3 thio C_{1-6} alkyl, Ar^3 aminocarbonyl, Het-aminocarbonyl, $Ar^3(CH_2)_n$ aminocarbonyl, Het- $(CH_2)_n$ aminocarbonyl; and the other one of R^{2a} and R^{2b} is hydrogen; or
 - 30 (d) one of R^{2a} and R^{3a} is selected from cyano C_{1-6} alkyl, Ar^3C_{1-6} alkyl, Het- C_{1-6} alkyl, $N(R^{8a}R^{8b})C_{1-6}$ alkyl, Ar^3C_{2-6} alkenyl, Ar^3 amino C_{1-6} alkyl, Het-amino C_{1-6} alkyl, Ar^3 aminocarbonyl; and the other one of R^{2a} and R^{2b} is hydrogen; or
 - 35 (e) one of R^{2a} and R^{3a} is selected from Ar^3C_{1-6} alkyl, $N(R^{8a}R^{8b})C_{1-6}$ alkyl, Ar^3C_{2-6} alkenyl, Ar^3 amino C_{1-6} alkyl; and the other one of R^{2a} and R^{2b} is hydrogen; or

(f) one of R^{2a} and R^{3a} is selected from $N(R^{8a}R^{8b})C_{1-6}alkyl$, $Ar^3aminoC_{1-6}alkyl$; and the other one of R^{2a} and R^{2b} is hydrogen;

and for any of (a) – (f)

in case R^{2a} is hydrogen then R^3 is hydrogen;

- 5 in case R^{2b} is hydrogen then R^3 is hydrogen or $C_{1-6}alkyl$; or preferably then R^3 is hydrogen.

Further particular subgroups of the compounds of formula (I) are those compounds of formula (I), or any subgroup of compounds of formula (I) specified herein, wherein R^5 is hydrogen.

Other particular subgroups of the compounds of formula (I) are those compounds of formula (I), or any subgroup of compounds of formula (I) specified herein, wherein Q is Ar^2 .

15 Other particular subgroups of the compounds of formula (I) are those compounds of formula (I), or any subgroup of compounds of formula (I) specified herein, wherein Q is R^{6a} .

20 Other particular subgroups of the compounds of formula (I) are those compounds of formula (I), or any subgroup of compounds of formula (I) specified herein, wherein Q is pyrrolidinyl substituted with R^6 , piperidinyl substituted with R^6 or homopiperidinyl substituted with R^6 ; in particular wherein Q is piperidinyl substituted with R^6 . Preferably the R^6 group is substituted on the nitrogen atom of the said pyrrolidinyl, piperidinyl or homopiperidinyl. More preferably the said pyrrolidinyl, piperidinyl or homopiperidinyl is linked to the $-N(R^5)-$ moiety via a 3-yl or in particular via a 4-yl link.

Interesting subgroups of compounds are those compounds of formula (I) or of any of the subgroups specified herein, wherein Q is R^{6a} , wherein

- (a) R^{6a} is $C_{1-6}alkyl$ substituted with two substituents or, preferably, with one substituent, each independently selected from the group consisting of trifluoromethyl, $NR^{7a}R^{7b}$, Ar^2 , hydroxy, $C_{1-4}alkoxy$, $Ar^2(CH_2)_n$ oxy, hydroxycarbonyl, aminocarbonyl, $C_{1-4}alkylcarbonyl$, $C_{1-4}alkoxycarbonyl$, $Ar^2(CH_2)_n$ carbonyl, aminocarbonyloxy, $C_{1-4}alkylcarbonyloxy$, Ar^2 carbonyloxy, mono- or di($C_{1-4}alkyl$)aminocarbonyl, aminosulfonyl, mono- or di($C_{1-4}alkyl$)aminosulfonyl or a heterocycle selected from the group consisting of pyrrolidinyl, imidazolyl, piperidinyl, homopiperidinyl, piperazinyl, dioxolanyl, dioxanyl and

- pyridyl, wherein each of said heterocycle may optionally be substituted with with one or two radicals selected from oxo and C₁₋₆alkyl; or in particular wherein
- (b) R^{6a} is C₁₋₆alkyl substituted with one substituent and, optionally with a further substituent which is hydroxy, wherein said substituent is trifluoromethyl, NR^{7a}R^{7b}, Ar², hydroxy, C₁₋₄alkoxy, Ar²(CH₂)_noxy, hydroxycarbonyl, aminocarbonyl, C₁₋₄alkylcarbonyl, C₁₋₄alkoxycarbonyl, Ar²(CH₂)_ncarbonyl, aminocarbonyloxy, C₁₋₄alkylcarbonyloxy, Ar²carbonyloxy, mono- or di(C₁₋₄alkyl)aminocarbonyl, aminosulfonyl, mono- or di(C₁₋₄alkyl)aminosulfonyl or a heterocycle selected from the group consisting of pyrrolidinyl, imidazolyl, piperidinyl, homopiperidinyl, piperazinyl, dioxolanyl, dioxanyl and pyridyl, wherein each of said heterocycle may optionally be substituted with with one or two radicals selected from oxo and C₁₋₆alkyl; or further in particular
- (c) R^{6a} is C₁₋₆alkyl substituted with NR^{7a}R^{7b}, Ar², hydroxy, C₁₋₄alkoxy, hydroxycarbonyl, aminocarbonyl, aminosulfonyl C₁₋₄alkylcarbonyl, Ar²carbonyl, C₁₋₄alkoxycarbonyl, or C₁₋₆alkyl substituted with two hydroxy radicals, or C₁₋₆alkyl substituted with a heterocycle selected from dioxolanyl, pyrrolidinyl, piperidinyl, homopiperidinyl, piperazinyl, wherein each of said heterocycle may optionally be substituted with oxo or with one or two C₁₋₆alkyl radicals; or further in particular
- (d) R^{6a} is C₁₋₆alkyl substituted with NR^{7a}R^{7b}, Ar², hydroxy, hydroxycarbonyl, aminocarbonyl, aminosulfonyl or C₁₋₆alkyl substituted with two hydroxy radicals, or C₁₋₆alkyl substituted with a heterocycle selected from dioxolanyl, pyrrolidinyl, piperidinyl, homopiperidinyl, piperazinyl, wherein each of said heterocycle may optionally be substituted with oxo or with one or two C₁₋₆alkyl radicals; or further in particular
- (e) R^{6a} is C₁₋₆alkyl substituted with Ar² or hydroxy, or C₁₋₆alkyl substituted with two hydroxy radicals, or C₁₋₆alkyl substituted with a heterocycle selected from dioxolanyl, pyrrolidinyl, piperidinyl, piperazinyl, wherein each of said heterocycle may optionally be substituted with one or two C₁₋₆alkyl radicals; or preferably
- (f) R^{6a} is C₁₋₆alkyl substituted with Ar² or hydroxy, or C₁₋₆alkyl substituted with two hydroxy radicals, or C₁₋₆alkyl substituted with diC₁₋₆alkyl-dioxolanyl, pyrrolidinyl, piperidinyl, piperazinyl, 4-C₁₋₆alkyl-piperazinyl; or preferably
- (g) R^{6a} is C₁₋₆alkyl substituted with Ar² or hydroxy, or C₁₋₆alkyl substituted with two hydroxy radicals, or C₁₋₆alkyl substituted with piperidinyl or with piperazinyl; or more preferably
- (h) R^{6a} is C₁₋₆alkyl substituted with Ar² or C₁₋₆alkyl substituted with piperidinyl or with piperazinyl.

Preferably in (a) – (h) in the previous paragraph the radicals pyrrolidinyl, piperidinyl, homopiperidinyl or piperazinyl are linked by their nitrogen atom to the C₁₋₆alkyl on which they are substituted.

- 5 Other particular subgroups of the compounds of formula (I) are those compounds of formula (I), or any subgroup of compounds of formula (I) specified herein, wherein Q is pyrrolidinyl substituted with R⁶, piperidinyl substituted with R⁶ or homopiperidinyl substituted with R⁶; wherein
 - 10 (a) R⁶ is hydrogen or C₁₋₆alkyl optionally substituted with two substituents or, preferably, with one substituent, each independently selected from the group consisting of trifluoromethyl, NR^{7a}R^{7b}, Ar², hydroxy, C₁₋₄alkoxy, Ar²(CH₂)_noxy, hydroxycarbonyl, aminocarbonyl, C₁₋₄alkylcarbonyl, C₁₋₄alkoxycarbonyl, Ar²(CH₂)_ncarbonyl, aminocarbonyloxy, C₁₋₄alkylcarbonyloxy, Ar²carbonyloxy, mono- or di(C₁₋₄alkyl)aminocarbonyl, aminosulfonyl, mono- or di(C₁₋₄alkyl)-
 - 15 aminosulfonyl or a heterocycle selected from the group consisting of pyrrolidinyl, imidazolyl, piperidinyl, homopiperidinyl, piperazinyl, dioxolanyl, dioxanyl and pyridyl, wherein each of said heterocycle may optionally be substituted with with one or two radicals selected from oxo and C₁₋₆alkyl; or in particular
 - 20 (b) R⁶ is hydrogen or C₁₋₆alkyl optionally substituted with one substituent and, optionally with a further substituent which is hydroxy, wherein said substituent is trifluoromethyl, NR^{7a}R^{7b}, Ar², hydroxy, C₁₋₄alkoxy, Ar²(CH₂)_noxy, hydroxycarbonyl, aminocarbonyl, C₁₋₄alkylcarbonyl, C₁₋₄alkoxycarbonyl, Ar²(CH₂)_ncarbonyl, aminocarbonyloxy, C₁₋₄alkylcarbonyloxy, Ar²carbonyloxy, mono- or di(C₁₋₄alkyl)aminocarbonyl, aminosulfonyl, mono- or di(C₁₋₄alkyl)aminosulfonyl
 - 25 or a heterocycle selected from the group consisting of pyrrolidinyl, imidazolyl, piperidinyl, homopiperidinyl, piperazinyl, dioxolanyl, dioxanyl and pyridyl, wherein each of said heterocycle may optionally be substituted with with one or two radicals selected from oxo and C₁₋₆alkyl; or further in particular
 - 30 (c) R⁶ is hydrogen or C₁₋₆alkyl optionally substituted with NR^{7a}R^{7b}, Ar², hydroxy, C₁₋₄alkoxy, hydroxycarbonyl, aminocarbonyl, aminosulfonyl, C₁₋₄alkoxycarbonyl or C₁₋₆alkyl substituted with two hydroxy radicals, or C₁₋₆alkyl substituted with a heterocycle selected from dioxolanyl, pyrrolidinyl, piperidinyl, homopiperidinyl, piperazinyl, wherein each of said heterocycle may optionally be substituted with oxo or with one or two C₁₋₆alkyl radicals; or further in particular
 - 35 (d) R⁶ is hydrogen or C₁₋₆alkyl optionally substituted with NR^{7a}R^{7b}, Ar², hydroxy, hydroxycarbonyl, aminocarbonyl, aminosulfonyl or C₁₋₆alkyl substituted with two hydroxy radicals, or C₁₋₆alkyl substituted with a heterocycle selected from dioxolanyl, pyrrolidinyl, piperidinyl, homopiperidinyl, piperazinyl, wherein each

of said heterocycle may optionally be substituted with oxo or with one or two C₁₋₆alkyl radicals; or further in particular wherein

- (e) R⁶ is hydrogen or C₁₋₆alkyl optionally substituted with Ar², hydroxy, aminocarbonyl or aminosulfonyl, or C₁₋₆alkyl substituted with two hydroxy radicals, or C₁₋₆alkyl substituted with a heterocycle selected from dioxolanyl, pyrrolidinyl, piperidinyl, piperazinyl, wherein each of said heterocycle may optionally be substituted with one or two C₁₋₆alkyl radicals; or preferably
- (f) R⁶ is hydrogen or C₁₋₆alkyl optionally substituted with Ar², hydroxy, aminocarbonyl, aminosulfonyl, or C₁₋₆alkyl substituted with two hydroxy radicals, or C₁₋₆alkyl substituted with pyrrolidinyl, piperidinyl, piperazinyl, 4-C₁₋₆alkylpiperazinyl; or preferably
- (g) R⁶ is hydrogen or C₁₋₆alkyl optionally substituted with Ar², hydroxy, aminocarbonyl or aminosulfonyl; or preferably
- (h) R⁶ is hydrogen or C₁₋₆alkyl substituted with Ar² or C₁₋₆alkyl substituted with piperidinyl or with piperazinyl; or
- (i) R⁶ is C₁₋₆alkyl.

Preferably in (a) – (h) in the previous paragraph, the radicals pyrrolidinyl, piperidinyl, homopiperidinyl or piperazinyl are linked by their nitrogen atom to the C₁₋₆alkyl on which they are substituted.

Further particular subgroups of the compounds of formula (I) are those compounds of formula (I), or any subgroup of compounds of formula (I) specified herein, wherein R^{7a} and R^{7b} are hydrogen or C₁₋₆alkyl, or preferably wherein R^{7a} and R^{7b} are hydrogen.

Further particular subgroups of the compounds of formula (I) are those compounds of formula (I), or any subgroup of compounds of formula (I) specified herein, wherein

- (a) R^{8a} is Ar³, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, C₁₋₆alkoxyC₁₋₆alkyl, cyanoC₁₋₆alkyl, aminoC₁₋₆alkyl, mono- or di(C₁₋₆alkyl)aminoC₁₋₆alkyl, Ar³C₁₋₆alkyl, Het-C₁₋₆alkyl, aminocarbonyl-C₁₋₆alkyl, carboxyl-C₁₋₆alkyl; and R^{8b} is Ar³; or
- (b) R^{8a} is Ar³, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, Ar³C₁₋₆alkyl, Het-C₁₋₆alkyl, aminocarbonyl-C₁₋₆alkyl; and R^{8b} is Ar³; or
- (c) R^{8a} is hydroxyC₁₋₆alkyl, aminocarbonyl-C₁₋₆alkyl; and R^{8b} is Ar³; or
- (d) R^{8a} is Ar³ and R^{8b} is Ar³; or
- (e) R^{8a} is C₁₋₆alkyl, hydroxyC₁₋₆alkyl, Ar³C₁₋₆alkyl, Het-C₁₋₆alkyl, aminocarbonyl-C₁₋₆alkyl; and R^{8b} is C₁₋₆alkyl, hydroxyC₁₋₆alkyl, Ar³C₁₋₆alkyl, Het-C₁₋₆alkyl.

In particular, Ar¹ is phenyl or phenyl substituted with 1, 2, 3 substituents or with 1, 2 substituents selected from those mentioned in the definition of the compounds of formula (I) or of any subgroup thereof.

- 5 In the group of compounds of formula (I) or in any of the subgroups of compounds of formula (I):
- (a) Ar¹ preferably is phenyl or phenyl substituted with up to 3 substituents, or with up to 2 substituents, or with one substituent, selected from halo, hydroxy, C₁₋₆alkyl, hydroxyc₁₋₆alkyl, trifluormethyl, and C₁₋₆alkyloxy;
 - 10 (b) Ar¹ more preferably is phenyl or phenyl substituted with up to 3 substituents, or with up to 2 substituents, or with one substituent, selected from halo, hydroxy, C₁₋₆alkyl and C₁₋₆alkyloxy;
 - (c) Ar¹ more preferably is phenyl or phenyl substituted with up to 3 substituents, or with up to 2 substituents, or with one substituent, selected from halo and
 - 15 C₁₋₆alkyl.

In particular, Ar² is phenyl or phenyl substituted with 1, 2, 3 substituents or with 1, 2 substituents selected from the group consisting of those mentioned in the definition of the compounds of formula (I) or of any subgroup thereof.

20

Further particular subgroups of the compounds of formula (I) are those compounds of formula (I), or any subgroup of compounds of formula (I) specified herein, wherein Ar² is as defined for Ar¹.

- 25 Further particular subgroups of the compounds of formula (I) are those compounds of formula (I), or any subgroup of compounds of formula (I) specified herein, wherein Ar³ is phenyl, naphthalenyl, 1,2,3,4-tetrahydro-naphthalenyl or indanyl, or preferably wherein Ar³ is phenyl, naphthalenyl or indanyl; wherein said phenyl may optionally and each individually be substituted with one or more, such as 2, 3 or 4, substituents
- 30 selected from the group consisting of substituents of Ar³ in the definitions of the compounds (I).

Further particular subgroups of the compounds of formula (I) are those compounds of formula (I), or any subgroup of compounds of formula (I) specified herein, wherein

- 35 (a) Ar³ is phenyl optionally substituted with one, two or three substituents selected from halo, hydroxy, mercapto, amino, cyano, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, Ar¹, hydroxyc₁₋₆alkyl, CF₃, aminoc₁₋₆alkyl, cyanoc₁₋₆alkyl, aminocarbonyl, C₁₋₆alkyloxy, C₁₋₆alkylthio, Ar¹-oxy, Ar¹-thio, Ar¹-amino, aminosulfonyl,


- aminocarbonyl-C₁₋₆alkyl, hydroxycarbonyl-C₁₋₆alkyl, hydroxycarbonyl, C₁₋₄alkylcarbonyl, C₁₋₄alkylcarbonylamino or C₁₋₄alkoxycarbonyl; or wherein
- (b) Ar³ is phenyl optionally substituted with one, two or three substituents selected from halo, hydroxy, mercapto, cyano, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, Ar¹, hydroxyC₁₋₆alkyl, CF₃, aminoC₁₋₆alkyl, cyanoC₁₋₆alkyl, aminocarbonyl, C₁₋₆alkyloxy, C₁₋₆alkylthio, Ar¹-oxy, Ar¹-thio, Ar¹-amino, aminosulfonyl, aminocarbonyl-C₁₋₆alkyl, hydroxycarbonyl-C₁₋₆alkyl, C₁₋₄alkylcarbonyl, C₁₋₄alkylcarbonylamino or C₁₋₄alkoxycarbonyl; or wherein
- (c) Ar³ is phenyl optionally substituted with one, two or three substituents selected from halo, C₁₋₆alkyl or hydroxyC₁₋₆alkyl; or
- (d) Ar³ is phenyl substituted with one, two or three substituents selected from halo, C₁₋₆alkyl or hydroxyC₁₋₆alkyl; or
- (e) Ar³ is phenyl optionally substituted with one, two or three substituents selected from halo, C₁₋₆alkyl or hydroxyC₁₋₆alkyl; or
- (f) Ar³ is phenyl substituted with one, two or three substituents selected from halo, C₁₋₆alkyl or hydroxyC₁₋₆alkyl; or
- (g) Ar³ is phenyl substituted with one or two substituents selected from halo, C₁₋₆alkyl or hydroxyC₁₋₆alkyl; or
- (h) Ar³ is phenyl optionally substituted with one or two substituents selected from C₁₋₆alkyl or hydroxyC₁₋₆alkyl.

In particular Ar³ is as defined for Ar², more in particular Ar³ is as defined for Ar¹.

- Further particular subgroups of the compounds of formula (I) are those compounds of formula (I), or any subgroup of compounds of formula (I) specified herein, wherein
- (a) Het is tetrahydrofuranyl, furanyl, thienyl, thiazolyl, oxazolyl, imidazolyl, isothiazolyl, pyrazolyl, isoxazolyl, piperidiny, homopiperidiny, piperaziny, morpholiny, pyridyl, pyraziny, pyrimidiny, tetrahydroquinoliny, quinoliny, isoquinoliny, benzodioxany, benzodioxoly, indoliny, indoly, which may optionally be substituted with oxo, amino, Ar¹, C₁₋₄alkyl, aminoC₁₋₄alkyl, hydroxyC₁₋₆alkyl, Ar¹C₁₋₄alkyl, mono- or di(C₁₋₆alkyl)aminoC₁₋₆alkyl, mono- or di(C₁₋₆alkyl)amino, (hydroxyC₁₋₆alkyl)amino, and optionally further with one or two C₁₋₄alkyl radicals; or
- (b) Het is tetrahydrofuranyl, furanyl, thienyl, thiazolyl, oxazolyl, imidazolyl, pyrazolyl, isoxazolyl, piperidiny, homopiperidiny, piperaziny, morpholiny, pyridyl, pyraziny, pyrimidiny, tetrahydroquinoliny, quinoliny, isoquinoliny, benzodioxany, benzodioxoly, indoliny, indoly, which may optionally be substituted with oxo, amino, Ar¹, C₁₋₄alkyl, aminoC₁₋₄alkyl, hydroxyC₁₋₆alkyl, and

- optionally further with one or two C₁₋₄alkyl radicals; or
- (c) Het is furanyl, thienyl, imidazolyl, pyrazolyl, isoxazolyl, morpholinyl, pyrimidinyl, benzodioxolyl, quinolinyl, indolinyl, which may optionally be substituted with hydroxyC₁₋₆alkyl or with one or two C₁₋₄alkyl radicals; or
- 5 (d) Het is furanyl, thienyl, imidazolyl, pyrazolyl, isoxazolyl, morpholinyl, pyrimidinyl, benzodioxolyl, quinolinyl, indolinyl, which may optionally be substituted with hydroxyC₁₋₆alkyl or with one or two C₁₋₄alkyl radicals; or
- (e) Het is furanyl, imidazolyl, morpholinyl, benzodioxolyl, quinolinyl, indolinyl, which may optionally be substituted with hydroxyC₁₋₆alkyl or with one or two
- 10 C₁₋₄alkyl radicals; or
- (f) Het is morpholinyl, which may optionally be substituted with one or two C₁₋₄alkyl radicals; or
- (g) Het is morpholinyl.
- 15 Preferred compounds are those compounds listed in tables 1 through 5, more in particular the compound numbers 1 to 77, 138, 143 to 165 and 171 to 177.


The compounds of formula (I) or any of the subgroups thereof can be prepared as in the following reaction schemes.



- In this schemes Q, G, R¹, R^{2a}, R^{2b}, R^{3a}, R^{3b}, R⁵ have the meanings defined above for the compounds of formula (I) or of any of the subgroups thereof. W is an appropriate leaving group, preferably it is chloro or bromo. The reaction of this scheme is typically conducted in a suitable solvent such as an ether, e.g. THF, a halogenated hydrocarbon, e.g. dichloromethane, CHCl₃, toluene, a polar aprotic solvent such as DMF, DMSO, DMA and the like.
- 25


- The compounds of formula (I) wherein Q is a pyrrolinyl, piperdinyl or homopiperidinyl group substituted with R⁶ which is other than hydrogen, i.e. R^{6a}, which are represented by formula (I-c-1), can be prepared as in the following reaction scheme.
- 30

-26-





- In this scheme G, t, R¹, R^{2a}, R^{2b}, R³, R⁵ have the meanings defined above for the compounds of formula (I) or of any of the subgroups thereof. W is an appropriate leaving group, preferably it is chloro or bromo. The reaction of this scheme can be conducted in a suitable solvent such as an ether, e.g. THF, a halogenated hydrocarbon, e.g. dichloromethane, CHCl₃, toluene, a polar aprotic solvent such as DMF, DMSO, DMA and the like. A base may be added to pick up the acid that is liberated during the reaction. If desired, certain catalysts such as iodide salts (e.g. KI) may be added.
- Intermediates (IV) can also be converted to compounds (I-c-1) with a reductive N-alkylation reaction starting from an aldehyde or ketone R^{6b}=O (V-a), wherein R^{6a} has the same meaning as R^{6a} provided that it has one hydrogen atom less. This reductive alkylation is done in a suitable solvent, e.g. an alcohol, using hydrogen in the presence of a metal catalyst such as Pd or NaBH₃CN.
- Some of the compounds of formula (I) can also be prepared starting from precursors of the compounds of formula (I) using appropriate functional group transformation reactions.
- Precursors of the compounds of formula (I) for example are those wherein R^{2a} or R^{2b} is C₁₋₆alkoxycarbonyl or C₁₋₆alkyl substituted with C₁₋₆alkoxycarbonyl, which can be reduced, e.g. with LiAlH₄, to the corresponding compounds of formula (I) wherein R^{2a} or R^{2b} is hydroxyC₁₋₆alkyl. The latter group can be oxidized with a mild oxidant to an aldehyde group, e.g. with MnO₂, which can further be derivatized with amines, e.g. with a reductive amination process, to the corresponding mono(derivatizedC₁₋₆alkyl)-amines. The latter can be alkylated or arylated to alkylamines wherein R^{2a} or R^{2b} has formula -Alk-NR^{8a}R^{8b}. Alternatively precursors of the compounds of formula (I) wherein R^{2a} or R^{2b} is hydroxyC₁₋₆alkyl can be converted to the corresponding haloC₁₋₆alkyl compounds, e.g. by treatment with a suitable halogenating agent such as SOCl₂, which compounds subsequently are reacted with an amine or amine derivative.
- This reaction sequence is illustrated by the following schemes in which R¹² represents a C₁₋₆alkyl radical, which preferably is methyl or ethyl. This reaction sequence may be done starting from (VII-a) or (VII-b) separately but also can be done using a mixture of

(VII-a) and (VII-b) and subsequently separating the reaction products either at the end of the reaction sequence or in an intermediate step.





5



Or starting from (VII-b):


10






- 5 Precursors of the compounds of formula (I) wherein R^{2a} or R^{2b} is an aldehyde can be converted to the corresponding compounds wherein R^{2a} or R^{2b} is substituted C_{2-6} alkenyl (I-g-1) or (I-g-2), by a Wittig reaction or a Wittig-Horner reaction. In the former instance a Wittig type reagent is used, such as a triphenylphosphoniumylide in a suitable reaction-inert solvent such as an ether, starting from triphenylphosphine and a
- 10 halo derivative. The Wittig-Horner reaction is performed using a phosphonate, such as e.g. a reagent of formula $\text{di}(C_{1-6}\text{alkyloxy})\text{-P(=O)-CH}_2\text{-CH}_2\text{-CN}$ in the presence of a base, preferably a strong base, in an aprotic organic solvent. Compounds wherein R^{2a} or R^{3a} is substituted C_{2-6} alkenyl can be reduced to the corresponding compounds wherein R^{2a} or R^{3a} is substituted C_{2-6} alkyl (I-i-1) or (I-i-2), e.g. with hydrogen in the presence of
- 15 a noble metal catalyst such as Pd/C. The cyano group in turn can be reduced to the corresponding methyleneamine ($-\text{CH}_2\text{-NH}_2$) group with hydrogen in the presence of a catalyst such as Raney Ni, in a suitable solvent such as methanol/ammonia. This reaction yields compounds (I-j-1) and (I-j-2) which can be mono-alkylated or double alkylated to yield compounds (I-k-1), (I-k-2) and (I-l-1), (I-l-2). These alkylations may
- 20 be done by a reductive alkylation reaction using an aldehyde or ketone in the presence of hydrogen and a catalyst (yielding mono-alkyl derivatives) or with suitably

- substituted alkyl halides (yielding mono- or dialkyl derivatives). These reactions are depicted in the following reaction schemes. In these schemes R^{2a-1} represents CN, Ar^3 or Het, Alk^1 represents C_{4-6} alkanediyl radicals (which are as C_{1-6} alkanediyl, but having from 4 – 6 carbon atoms), R^{8a} and R^{8b} have the same meanings as defined in this specification and claims, but preferably are other than Ar^3 .
- 5



-30-



5

Compounds of formula (I) wherein R^{2a} or R^{2b} is an aldehyde or C₁₋₆alkyl substituted with a keto or an aldehyde can also be derivatized with a Grignard type of reaction to introduce aryl or alkyl groups.

An additional aspect of the present invention concerns the fact that some of the compounds identified as precursors of the compounds of formula (I), are novel compounds.

- 5 In particular the compounds of formula (VII-a), (VII-b), (VIII-a), (VIII-b), (IX-a), (IX-b), (I-f-1), (I-f-2), (I-g-1), (I-g-2) wherein G , R^1 , R^{2a} , R^{2b} , R^3 , R^5 , R^{8a} , R^{8b} , R^{12} are as defined above in the definitions of the compounds of formula (I) or in any of the subgroups thereof, and wherein Q is pyrrolidinyl, piperidinyl or homopiperidinyl, substituted on their nitrogen with a radical R^6 which is C_{1-6} alkyl optionally substituted
- 10 with one or more, preferably one or two, substituents each independently selected from the group consisting of trifluoromethyl, C_{3-7} cycloalkyl, Ar^2 , hydroxy, C_{1-4} alkoxy, C_{1-4} alkylthio, Ar^2 -oxy-, Ar^2 -thio-, $Ar^2(CH_2)_n$ oxy, $Ar^2(CH_2)_n$ thio, hydroxycarbonyl, aminocarbonyl, C_{1-4} alkylcarbonyl, Ar^2 carbonyl, C_{1-4} alkoxycarbonyl, $Ar^2(CH_2)_n$ carbonyl, aminocarbonyloxy, C_{1-4} alkylcarbonyloxy, Ar^2 carbonyloxy,
- 15 $Ar^2(CH_2)_n$ carbonyloxy, C_{1-4} alkoxycarbonyl $(CH_2)_n$ oxy, mono- or di(C_{1-4} alkyl)amino-carbonyl, mono- or di(C_{1-4} alkyl)aminocarbonyloxy, aminosulfonyl, mono- or di(C_{1-4} alkyl)aminosulfonyl or a heterocycle selected from the group consisting of pyrrolidinyl, pyrrolyl, dihydropyrrolyl, imidazolyl, triazolyl, piperidinyl, homopiperidinyl, piperazinyl, dioxolanyl, dioxanyl, pyridyl and tetrahydropyridyl,
- 20 wherein each of said heterocycle may optionally be substituted with one or two substituents selected from oxo or C_{1-6} alkyl; and wherein said R^6 can be represented by R^{6b} , as well as the pharmaceutically acceptable salt forms thereof, and the possible stereoisomeric forms thereof, are novel compounds.
- 25 Of particular interest are any of the groups of novel compounds specified in the previous paragraph wherein:
- (a) R^{6b} is C_{1-6} alkyl optionally substituted with two substituents or, preferably, with one substituent, each independently selected from the group consisting of trifluoromethyl, Ar^2 , hydroxy, C_{1-4} alkoxy, $Ar^2(CH_2)_n$ oxy, hydroxycarbonyl,
- 30 aminocarbonyl, C_{1-4} alkylcarbonyl, C_{1-4} alkoxycarbonyl, $Ar^2(CH_2)_n$ carbonyl, aminocarbonyloxy, C_{1-4} alkylcarbonyloxy, Ar^2 carbonyloxy, mono- or di(C_{1-4} alkyl)amino-carbonyl, aminosulfonyl, mono- or di(C_{1-4} alkyl)aminosulfonyl or a heterocycle selected from the group consisting of pyrrolidinyl, imidazolyl, piperidinyl, homopiperidinyl, piperazinyl, dioxolanyl, dioxanyl and pyridyl, wherein each of
- 35 said heterocycle may optionally be substituted with with one or two radicals selected from oxo and C_{1-6} alkyl; or in particular
- (b) R^{6b} is C_{1-6} alkyl optionally substituted with one substituent and, optionally with a further substituent which is hydroxy, wherein said substituent is trifluoromethyl,

- Ar², hydroxy, C₁₋₄alkoxy, Ar²(CH₂)_noxy, hydroxycarbonyl, aminocarbonyl, C₁₋₄alkylcarbonyl, C₁₋₄alkoxycarbonyl, Ar²(CH₂)_ncarbonyl, aminocarbonyloxy, C₁₋₄alkylcarbonyloxy, Ar²carbonyloxy, mono- or di(C₁₋₄alkyl)aminocarbonyl, aminosulfonyl, mono- or di(C₁₋₄alkyl)aminosulfonyl or a heterocycle selected from the group consisting of pyrrolidinyl, imidazolyl, piperidinyl, homopiperidinyl, piperazinyl, dioxolanyl, dioxanyl and pyridyl, wherein each of said heterocycle may optionally be substituted with one or two radicals selected from oxo and C₁₋₆alkyl; or further in particular
- (c) R^{6b} is C₁₋₆alkyl optionally substituted with Ar², hydroxy, C₁₋₄alkoxy, hydroxycarbonyl, aminocarbonyl, aminosulfonyl, C₁₋₄alkoxycarbonyl or C₁₋₆alkyl substituted with two hydroxy radicals, or C₁₋₆alkyl substituted with a heterocycle selected from dioxolanyl, pyrrolidinyl, piperidinyl, homopiperidinyl, piperazinyl, wherein each of said heterocycle may optionally be substituted with oxo or with one or two C₁₋₆alkyl radicals; or further in particular
- (d) R^{6b} is C₁₋₆alkyl optionally substituted with Ar², hydroxy, hydroxycarbonyl, aminocarbonyl, aminosulfonyl or C₁₋₆alkyl substituted with two hydroxy radicals, or C₁₋₆alkyl substituted with a heterocycle selected from dioxolanyl, pyrrolidinyl, piperidinyl, homopiperidinyl, piperazinyl, wherein each of said heterocycle may optionally be substituted with oxo or with one or two C₁₋₆alkyl radicals; or further in particular wherein
- (e) R^{6b} is C₁₋₆alkyl optionally substituted with Ar², hydroxy, aminocarbonyl or aminosulfonyl, or C₁₋₆alkyl substituted with two hydroxy radicals, or C₁₋₆alkyl substituted with a heterocycle selected from dioxolanyl, pyrrolidinyl, piperidinyl, piperazinyl, wherein each of said heterocycle may optionally be substituted with one or two C₁₋₆alkyl radicals; or preferably
- (f) R^{6b} is C₁₋₆alkyl optionally substituted with Ar², hydroxy, aminocarbonyl, aminosulfonyl, or C₁₋₆alkyl substituted with two hydroxy radicals, or C₁₋₆alkyl substituted with pyrrolidinyl, piperidinyl, piperazinyl, 4-C₁₋₆alkyl-piperazinyl; or preferably
- (g) R^{6b} is C₁₋₆alkyl optionally substituted with Ar², hydroxy, aminocarbonyl or aminosulfonyl; or preferably
- (h) R^{6b} is C₁₋₆alkyl.

Preferably in (a) — (h) in the previous paragraph the radicals pyrrolidinyl, piperidinyl, homopiperidinyl or piperazinyl are linked by their nitrogen atom to the C₁₋₆alkyl on which they are substituted.

Moreover these compounds have found to possess antiviral properties, in particular to possess RSV inhibitory activity. Of particular interest are the compounds of formula (VII-a), (VII-b), (VIII-a) and (VIII-b) wherein Q is 4-piperidinyl wherein the ring nitrogen is substituted with a radical R^6 which is C_{1-6} alkyl, as well as the
5 pharmaceutically acceptable salt forms thereof, and the possible stereoisomeric forms thereof. As used herein, the terms 'pharmaceutically acceptable salt forms' and the 'stereoisomeric forms' have the meanings specified above in this specification.

Also the compounds of formula (VII-a), (VII-b), (VIII-a), (VIII-b), (IX-a),
10 (IX-b), (I-f-1), (I-f-2), (I-g-1), (I-g-2) wherein G, R^1 , R^{2a} , R^{2b} , R^3 , R^5 , R^{8a} , R^{8b} and R^{12} are as defined above in the definitions of the compounds of formula (I) or in any of the subgroups thereof, and wherein Q is R^{6b} which is as specified in the previous paragraphs, as well as the pharmaceutically acceptable salt forms thereof, and the
15 possible stereoisomeric forms thereof, are novel compounds. Moreover these compounds have found to possess antiviral properties, in particular to possess RSV inhibitory activity.


Of particular interest are those compounds mentioned in the previous paragraph wherein G, R^1 , R^{2a} , R^{2b} , R^3 , R^5 , R^{8a} and R^{8b} are as specified in any of the subgroups
20 mentioned in this specification and claims. Preferred are those novel compounds mentioned in the previous paragraphs wherein G is C_{1-6} alkanediyl, more preferably wherein G is methylene; and/or wherein R^{2a} , R^{2b} , R^3 , R^5 are all hydrogen; and/or R^1 is pyridyl being substituted as outlined in this specification and claims, in particular R^1 is pyridyl being substituted with one or two substituents selected from C_{1-6} alkyl and
25 hydroxy.

Compounds of formula (I) may be converted into each other following art-known functional group transformation reactions, comprising those described hereinafter.

30 Nitro groups can be reduced to amino groups, which subsequently may be alkylated to mono- or dialkylamino groups, or acylated to arylcarbonylamino or alkylcarbonylamino and the like groups. Cyano groups may be reduced to aminomethylene groups, which similarly may be derivatized.

35 A number of the intermediates used to prepare the compounds of formula (I) are known compounds or are analogs of known compounds, which can be prepared following modifications of art-known methodologies readily accessible to the skilled person. A number of preparations of intermediates are given hereafter in somewhat more detail.

-34-



In a first step, a diaminobenzene (VI) is cyclized with urea in a suitable solvent, e.g. xylene, to yield a benzimidazolone (VII). The latter is converted to a benzimidazole derivative (VIII) wherein W is a leaving group as specified above, in particular by
 5 reaction of (VII) with a suitable halogenating agent, for example POCl_3 , and the resulting intermediate (VIII) is reacted with the amine derivative (IX) to obtain intermediate (II).

The same reaction sequence may be used to prepare other intermediates. For example
 10 the intermediates of formula (IV) can be prepared by reacting intermediates (IX) with an amine (X) wherein Q is a pyrrolinyl, piperidinyl or homopiperidinyl group wherein the nitrogen is substituted with a protective group to yield precursors of (IV) which can be converted to intermediates (IV) by removing the protective group. Suitable
 protecting groups for this purpose comprise alkyloxycarbonyl groups such as methoxy
 15 or ethoxycarbonyl, which can be removed with a base, or benzyl or benzyloxycarbonyl groups, which can be removed with hydrogen in the presence of a catalyst.

The compounds of formula (I) may be converted to the corresponding *N*-oxide forms following art-known procedures for converting a trivalent nitrogen into its *N*-oxide
 20 form. Said *N*-oxidation reaction may generally be carried out by reacting the starting material of formula (I) with an appropriate organic or inorganic peroxide. Appropriate inorganic peroxides comprise, for example, hydrogen peroxide, alkali metal or earth
 alkaline metal peroxides, e.g. sodium peroxide, potassium peroxide; appropriate
 organic peroxides may comprise peroxy acids such as, for example, benzenecarbo-
 25 peroxy acid or halo substituted benzenecarbo-
 peroxy acid, e.g. 3-chlorobenzenecarbo-
 peroxy acid, peroxyalkanoic acids, e.g. peroxyacetic acid, alkylhydroperoxides, e.g.
 t.butyl hydro-peroxide. Suitable solvents are, for example, water, lower alcohols, e.g.

ethanol and the like, hydrocarbons, e.g. toluene, ketones, e.g. 2-butanone, halogenated hydrocarbons, e.g. dichloromethane, and mixtures of such solvents.

5 Pure stereochemically isomeric forms of the compounds of formula (I) may be obtained by the application of art-known procedures. Diastereomers may be separated by physical methods such as selective crystallization and chromatographic techniques, e.g., counter-current distribution, liquid chromatography and the like.

10 The compounds of formula (I) as prepared in the hereinabove described processes are generally racemic mixtures of enantiomers which can be separated from one another following art-known resolution procedures. The racemic compounds of formula (I), which are sufficiently basic or acidic may be converted into the corresponding diastereomeric salt forms by reaction with a suitable chiral acid, respectively chiral base. Said diastereomeric salt forms are subsequently separated, for example, by selective or fractional crystallization and the enantiomers are liberated therefrom by alkali or acid. An alternative manner of
15 separating the enantiomeric forms of the compounds of formula (I) involves liquid chromatography, in particular liquid chromatography using a chiral stationary phase. Said pure stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting materials, provided that the
20 reaction occurs stereospecifically. Preferably if a specific stereoisomer is desired, said compound will be synthesized by stereospecific methods of preparation. These methods will advantageously employ enantiomerically pure starting materials.

In a further aspect, the present invention concerns a pharmaceutical composition
25 comprising a therapeutically effective amount of a compound of formula (I) as specified herein, or a compound of any of the subgroups of compounds of formula (I) as specified herein, and a pharmaceutically acceptable carrier. A therapeutically effective amount in this context is an amount sufficient to prophylactically act against, to stabilize or to reduce viral infection, and in particular RSV viral infection, in
30 infected subjects or subjects being at risk of being infected. In still a further aspect, this invention relates to a process of preparing a pharmaceutical composition as specified herein, which comprises intimately mixing a pharmaceutically acceptable carrier with a therapeutically effective amount of a compound of formula (I), as specified herein, or of a compound of any of the subgroups of compounds of formula (I) as specified
35 herein.

Therefore, the compounds of the present invention or any subgroup thereof may be formulated into various pharmaceutical forms for administration purposes. As

appropriate compositions there may be cited all compositions usually employed for systemically administering drugs. To prepare the pharmaceutical compositions of this invention, an effective amount of the particular compound, optionally in addition salt form or metal complex, as the active ingredient is combined in intimate admixture with

5 a pharmaceutically acceptable carrier, which carrier may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirable in unitary dosage form suitable, particularly, for administration orally, rectally, percutaneously, or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical

10 media may be employed such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs, emulsions and solutions; or solid carriers such as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules, and tablets. Because of their ease in administration, tablets and capsules represent the most

15 advantageous oral dosage unit forms, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and

20 glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations. In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a

25 suitable wetting agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not introduce a significant deleterious effect on the skin.

The compounds of the present invention may also be administered via oral inhalation or

30 insufflation by means of methods and formulations employed in the art for administration via this way. Thus, in general the compounds of the present invention may be administered to the lungs in the form of a solution, a suspension or a dry powder, a solution being preferred. Any system developed for the delivery of solutions, suspensions or dry powders via oral inhalation or insufflation are suitable for

35 the administration of the present compounds.

Thus, the present invention also provides a pharmaceutical composition adapted for administration by inhalation or insufflation through the mouth comprising a compound

of formula (I) and a pharmaceutically acceptable carrier. Preferably, the compounds of the present invention are administered via inhalation of a solution in nebulized or aerosolized doses.

- 5 It is especially advantageous to formulate the aforementioned pharmaceutical compositions in unit dosage form for ease of administration and uniformity of dosage. Unit dosage form as used herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required
10 pharmaceutical carrier. Examples of such unit dosage forms are tablets (including scored or coated tablets), capsules, pills, suppositories, powder packets, wafers, injectable solutions or suspensions and the like, and segregated multiples thereof.

- The compounds of formula (I) show antiviral properties. Viral infections treatable
15 using the compounds and methods of the present invention include those infections brought on by ortho- and paramyxoviruses and in particular by human and bovine respiratory syncytial virus (RSV). A number of the compounds of this invention moreover are active against mutated strains of RSV. Additionally, many of the compounds of this invention show a favorable pharmacokinetic profile and have
20 attractive properties in terms of bioavailability, including an acceptable half-life, AUC and peak values and lacking unfavourable phenomena such as insufficient quick onset and tissue retention.

- The *in vitro* antiviral activity against RSV of the present compounds was tested in a test
25 as described in the experimental part of the description, and may also be demonstrated in a virus yield reduction assay. The *in vivo* antiviral activity against RSV of the present compounds may be demonstrated in a test model using cotton rats as described in Wyde et al. (Antiviral Research (1998), 38, 31-42).

- 30 Due to their antiviral properties, particularly their anti-RSV properties, the compounds of formula (I) or any subgroup thereof, their prodrugs, *N*-oxides, addition salts, quaternary amines, metal complexes and stereochemically isomeric forms, are useful in the treatment of individuals experiencing a viral infection, particularly a RSV infection, and for the prophylaxis of these infections. In general, the compounds of the present
35 invention may be useful in the treatment of warm-blooded animals infected with viruses, in particular the respiratory syncytial virus.

The compounds of the present invention or any subgroup thereof may therefore be used as medicines. Said use as a medicine or method of treatment comprises the systemic

administration to viral infected subjects or to subjects susceptible to viral infections of an amount effective to combat the conditions associated with the viral infection, in particular the RSV infection.

- 5 The present invention also relates to the use of the present compounds or any subgroup thereof in the manufacture of a medicament for the treatment or the prevention of viral infections, particularly RSV infection.

- 10 The present invention furthermore relates to a method of treating a warm-blooded animal infected by a virus, or being at risk of infection by a virus, in particular by RSV, said method comprising the administration of an anti-virally effective amount of a compound of formula (I), as specified herein, or of a compound of any of the subgroups of compounds of formula (I), as specified herein.

- 15 In general it is contemplated that an antiviral effective daily amount would be from 0.01 mg/kg to 500 mg/kg body weight, more preferably from 0.1 mg/kg to 50 mg/kg body weight. It may be appropriate to administer the required dose as two, three, four or more sub-doses at appropriate intervals throughout the day. Said sub-doses may be formulated as unit dosage forms, for example, containing 1 to 1000 mg, and in
20 particular 5 to 200 mg of active ingredient per unit dosage form.

- The exact dosage and frequency of administration depends on the particular compound of formula (I) used, the particular condition being treated, the severity of the condition being treated, the age, weight, sex, extent of disorder and general physical condition of
25 the particular patient as well as other medication the individual may be taking, as is well known to those skilled in the art. Furthermore, it is evident that said effective daily amount may be lowered or increased depending on the response of the treated subject and/or depending on the evaluation of the physician prescribing the compounds of the instant invention. The effective daily amount ranges mentioned hereinabove are
30 therefore only guidelines.

- Also, the combination of another antiviral agent and a compound of formula (I) can be used as a medicine. Thus, the present invention also relates to a product containing (a) a compound of formula (I), and (b) another antiviral compound, as a combined
35 preparation for simultaneous, separate or sequential use in antiviral treatment. The different drugs may be combined in a single preparation together with pharmaceutically acceptable carriers. For instance, the compounds of the present invention may be combined with interferon-beta or tumor necrosis factor-alpha in order to treat or prevent RSV infections.

Examples


The following examples are intended to illustrate the present invention and not to limit it thereto. The terms 'compound 1', 'compound 2', etc. used in these examples refers to the same compounds in the tables.

5

The compounds were identified by LC/MS using the following equipment:


LCT: electrospray ionisation in positive mode, scanning mode from 100 to 900 amu; Xterra MS C18 (Waters, Milford, MA) 5 μ m, 3.9 x 150 mm); flow rate 1 ml/min. Two mobile phases (mobile phase A: 85% 6.5mM ammonium acetate + 15% acetonitrile; mobile phase B: 20% 6.5 mM ammonium acetate + 80% acetonitrile) were employed to run a gradient from 100 % A for 3 min to 100% B in 5 min., 100% B for 6 min to 100 % A in 3 min, and equilibrate again with 100 % A for 3 min).

ZQ: electrospray ionisation in both positive and negative (pulsed) mode scanning from 100 to 1000 amu; Xterra RP C18 (Waters, Milford, MA) 5 μ m, 3.9 x 150 mm); flow rate 1 ml/min. Two mobile phases (mobile phase A: 85% 6.5mM ammonium acetate + 15% acetonitrile; mobile phase B: 20% 6.5 mM ammonium acetate + 80% acetonitrile) were employed to run a gradient condition from 100 % A for 3 min to 100% B in 5 min., 100% B for 6 min to 100 % A in 3 min, and equilibrate again with 100 % A for 3 min).

Example 1Scheme A

25

-40-



- 5 A mixture of **a-1** (0.166 mol) and urea (0.199 mol) in xylene (300 ml) was stirred under reflux for 12 hours. The reaction was cooled down to room temperature. The precipitate was filtered off, rinsed with xylene and diisopropyl ether, and then dried, yielding 32g of intermediate **a-2** (93%, melting point: $> 260^\circ\text{C}$).
- 10 A mixture of **a-2** (0.073 mol) in POCl_3 (150 ml) was stirred at 100°C . HCl conc. (around 1.5 ml) was added drop wise very carefully until the dissolution of **a-2**. The mixture was stirred at 120°C for 6 hours. The solvent was evaporated until dryness. The residue was taken-up in $\text{H}_2\text{O}/\text{ice}$, basified with K_2CO_3 (powder) and extracted with ethylacetate + 10% methanol. The organic layer was separated, dried (over MgSO_4),
- 15 filtered and the solvent was evaporated until dryness, yielding 13.5 g of intermediate **a-3** (83%, melting point: 178°C).

A mixture of **a-3** (0.051 mmol) and **a-4** (0.056 mol) was stirred at 160°C for 2 hours. The residue was taken-up in $\text{CH}_2\text{Cl}_2/\text{H}_2\text{O}$ and basified with a 10% solution of K_2CO_3 in water. The organic layer was separated, dried (over MgSO_4), filtered and the solvent

20

was evaporated until dryness. The residue was purified by column chromatography over silica gel (eluent: CH_2Cl_2 /methanol/ NH_4OH 95/5/0.5). The pure fractions were collected and the solvent was evaporated, yielding 15.3 g of intermediate **a-5** (79%).

- 5 A mixture of **a-5** (0.0396 mol), **a-6** (0.059 mol) and K_2CO_3 (0.1584 mol) in CH_3CN (180ml) was stirred and refluxed for 12 hours. The solvent was evaporated until dryness. The residue was taken up in CH_2Cl_2 . The organic layer was washed with H_2O , dried (over MgSO_4), filtered and the solvent was evaporated until dryness. The residue (20g) was purified by column chromatography over silica gel (eluent: Toluene/
10 2-propanol/ NH_4OH 85/15/1; 20-45 μm). Two fractions were collected and the solvent was evaporated, yielding 5.3g of fraction 1 (27%) and 6.3g of fraction 2 (32%). Fraction 1 was crystallized twice in 2-propanone/ CH_3CN /diisopropylether. The precipitate was filtered off and dried, yielding 4.9g of intermediate **a-7** (25%, melting point: 179°C). Fraction 2 was crystallized from 2-propanone/ CH_3CN /diisopropylether.
15 The precipitate was filtered off and dried, yielding 5.5g of intermediate **a-8** (28%, melting point: 238°C).

- LiAlH_4 (0.009 mol) was added portion wise to a mixture of **a-7** (0.003 mol) in tetrahydrofuran (60 ml) at 5°C under N_2 flow. The reaction was stirred at 5°C for 1
20 hour and then at room temperature for 12 hours. Ethyl acetate and H_2O were added carefully and the aqueous layer was saturated with K_2CO_3 (powder). The organic layer was separated, dried (over MgSO_4) and then filtered over celite. The filtrate was evaporated until dryness, yielding 1.3 g of intermediate **a-9** (97%). The crude product was used directly in the next reaction step.

- 25 Intermediate **a-10** was prepared analogously to the procedure described for intermediate **a-9**.

- A mixture of **a-9** (0.0028 mol) and Pd/C 10% (2.5g) in CH_3OH (40ml) was hydrogenated at 40°C for 12 hours under an 8 bar pressure, then filtered over celite. Celite was washed with a solution of CH_3OH /tetrahydrofuran (50/50). The filtrate was
30 evaporated until dryness, yielding 1.8g of intermediate **a-11** (95%, melting point: 260°C).

- HCHO 37% in water (0.0098 mol), NaBH_3CN (0.0059 mol) then $\text{CH}_3\text{CO}_2\text{H}$ (2ml) were added at room temperature to a mixture of **a-11** (0.0049 mol) in CH_3CN (50ml). The
35 mixture was stirred at room temperature for 12 hours. The solvent was evaporated until dryness. The residue was taken up in ethanol (30ml) and a 5N solution of HCl in 2-propanol (4ml) was added. The mixture was stirred at 80°C for 8 hours. The solvent was evaporated until dryness. The residue was taken up in CH_2Cl_2 / K_2CO_3 10%. The

organic layer was separated, dried (over MgSO_4), filtered and the solvent was evaporated until dryness. The residue was crystallized from $\text{CH}_3\text{OH}/2$ -propanone/ CH_3CN . The precipitate was filtered off and dried, yielding 1.65g of **a-12** (88%). Part of this fraction (0.15g) was crystallized from $\text{CH}_3\text{OH}/2$ -propanone. The precipitate was

5 filtered off and dried (melting point: 165°C).

SOCl_2 (2.1ml) was added drop wise to a mixture of **a-12** (0.0018 mol) in CH_2Cl_2 (20ml) at 5°C . The mixture was stirred at 5°C for 1 hour, then at room temperature for 12 hours. The solvent was evaporated until dryness, yielding 0.93g of intermediate **a-13**

10 (100%). The crude product was used directly in the next reaction step.


A mixture of **a-13** (0.0003 mol), **a-14** (0.0005 mol) and K_2CO_3 (0.0019 mol) in dimethylformamide (30ml) was stirred at 80°C for 4 hours and poured into H_2O . The aqueous layer was saturated with K_2CO_3 and extracted with $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$. The

15 organic layer was separated, dried (over MgSO_4), filtered and the solvent was evaporated until dryness. The residue (0.25g) was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$ 90/10/1; $10\mu\text{m}$). The pure fractions were collected and the solvent was evaporated. The residue (0.05g, 24%) was

20 crystallized from 2-propanone/diisopropylether. The precipitate was filtered off and dried, yielding 0.042g of compound **a-15** (20%, compound **1**, melting point: 201°C).

Example 2

Scheme B



25 A mixture of **b-1** (0.0028 mol) and MnO_2 (2.5g) in CH_2Cl_2 (40ml) was stirred at room temperature for 12 hours, and then filtered over celite. Celite was rinsed with CH_2Cl_2 . The filtrate was evaporated until dryness. The residue was taken up in 2-propanone.


The precipitate was filtered off and dried, yielding 0.75g of intermediate **b-2** (69%, melting point: 250°C).

Variant 1 : A mixture of **b-2** (0.0001 mol), 3,5-dichloro aniline (0.0001 mol), BH_3CN on solid support (0.0001 mol) and $\text{CH}_3\text{CO}_2\text{H}$ (2 drops) in CH_3OH (4ml) was stirred at room temperature for 24 hours. The solution was filtered. The filtrate was evaporated until dryness. The residue was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$ 87/12/1.5; 5 μm). The pure fractions were collected and the solvent was evaporated, yielding 0.026g of 2-[6-[(3,5-dichloro-phenylamino)-methyl]-2-(1-methyl-piperidin-4-ylamino)-benzimidazol-1-ylmethyl]-6-methyl-pyridin-3-ol (38%, compound 2).

Variant 2 : **b-2** (0.0005 mol), NaBH_3CN (0.0006 mol), and then $\text{CH}_3\text{CO}_2\text{H}$ (0.2ml) were added at room temperature to a mixture of 3-methyl-aniline (0.0006 mol) in CH_3CN (20ml). The mixture was stirred at room temperature for 12 hours. H_2O was added. The mixture was saturated with K_2CO_3 (powder) and extracted with $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$. The organic layer was separated, dried (over MgSO_4), filtered and the solvent was evaporated until dryness. The residue (0.3g) was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{triethylamine}$; 90/10/0.1; 5 μm). The pure fractions were collected and the solvent was evaporated. The residue (0.17g, 68%) was crystallized from $\text{CH}_3\text{OH}/2$ -propanone/diisopropylether. The precipitate was filtered off and dried, yielding 0.13g of 2-[6-[(1-ethylidene-3-methyl-penta-2,4-dienylamino)-methyl]-2-(1-methyl-piperidin-4-ylamino)-benzimidazol-1-ylmethyl]-6-methyl-pyridin-3-ol (52%, compound 9, melting point: 141°C).

Example 3

Scheme C




A mixture of **c-1** (0.0018 mol), 2-methyl-thiophenol (0.002 mol) and K_2CO_3 (0.0077 mol) in CH_3CN (70ml) was stirred at 50°C for 12 hours. The solvent was evaporated until dryness. The residue was taken up in H_2O . The mixture was extracted with CH_2Cl_2 . The organic layer was separated, dried (over MgSO_4), filtered and the

solvent was evaporated until dryness. The residue (0.55g) was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$ 88/12/2; 15–40 μm). The pure fractions were collected and the solvent was evaporated. The residue (0.35g, 39%) was crystallized from $\text{CH}_3\text{CN}/\text{diisopropylether}$. The precipitate was filtered off and dried, yielding 0.32g of 6-methyl-2-[2-(1-methyl-piperidin-4-ylamino)-6-m-tolylsulfanylmethyl-benzoimidazol-1-ylmethyl]-pyridin-3-ol (compound 93, melting point: 202°C).

Example 4

10

Scheme D



In scheme D, R is defined as Ar^3 , Het^1 , $\text{Het}^1(\text{CH}_2)_n$ or $\text{Het}^1(\text{CH}_2)_n$.

15 Intermediate **d-2** (melting point: 262°C) was prepared analogous to the procedure described for intermediate **a-11**. Intermediate **d-3** was prepared analogous to the procedure described for intermediate **a-12**.


20 A mixture of **d-3** (0.0003 mol) in a 3N solution of HCl in water (30ml) was stirred at 80°C for 12 hours. The solvent was evaporated. The residue was dried, yielding 0.18g of intermediate **d-4**. The crude product was used directly in the next reaction step.

25 A mixture of **d-4** (0.0003 mol), 2-methyl-aniline (0.0005 mol), 1-(3-dimethylamino-propyl)-3-ethylcarbodiimide hydrochloride (0.0005 mol) and 1-hydroxybenzotriazole (0.0005 mol) in CH_2Cl_2 (20ml) was stirred at room temperature for 24 hours. A 10% solution of K_2CO_3 in water was added. The aqueous layer was saturated with K_2CO_3 (powder). The organic layer was separated, dried (over MgSO_4), filtered and the solvent was evaporated until dryness. The residue (0.2g) was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$ 80/20/1; 10 μm). The


pure fractions were collected and the solvent was evaporated. Yielding: 0.29g (14%). This fraction was taken up in diisopropyl ether, then CH₃OH/diisopropylether. The precipitate was filtered off and dried, yielding 0.007g of 3-(3-hydroxy-6-methyl-pyridin-2-ylmethyl)-2-(1-methyl-piperidin-4-ylamino)-3H-benzoimidazole-5-carboxylic acid m-tolylamide (4%, compound 116, melting point: 172°C).

Example 5

Scheme E



-46-



A mixture of **e-1** (0.0524 mol) and **e-2** (0.1048 mol) was stirred at 120°C in a Parr pressure vessel for 10 hours, then taken up into H₂O and extracted with ethyl acetate. The separated organic layer was purified by short open column chromatography over silica gel (eluent: CH₂Cl₂/methanol 96/4). The product fractions were collected and the solvent was evaporated until dryness, yielding 7.7g of intermediate **e-3** (44%).

A mixture of **e-3** (0.0312 mol), **e-4** (0.0343 mol) and K₂CO₃ (0.1092 mol) in dimethyl formamide (100ml) was stirred at 70°C for 24 hours. H₂O was then added. The mixture was extracted with CH₂Cl₂. The organic layer was separated, dried (over MgSO₄), filtered and the solvent was evaporated until dryness. The residue (12.2g) was purified by column chromatography over silica gel (eluent: toluene/isopropanol/NH₄OH 90/10/0.5; 15-40μm). Two fractions were collected and the solvent was evaporated, yielding 4g of intermediate **e-5** (28%) and 5.4g of intermediate **e-6** (38%).

e-5 (0.0088 mol) was added portion wise to a 48% solution of HBr in water (40ml). The mixture was brought slowly to 70°C, and then stirred for 12 hours. The precipitate was filtered, washed with CH₃CN and dried. The residue (4.6g, 80%) was taken up in H₂O and basified with K₂CO₃ (powder). The precipitate was filtered, and then washed with ethanol. The filtrate was evaporated, yielding 3g of intermediate **e-7** (52%). In an analogous way, **e-8** was prepared

HCHO 37% in H₂O (0.0152 mol) then NaBH₃CN (0.0091 mol) were added at room temperature to a mixture of **e-7** (0.0075 mol) in CH₃CN (100ml). Acetic acid (3.5ml) was added slowly at room temperature. The mixture was stirred at room temperature for 12 hours and poured into H₂O. The aqueous layer was saturated with K₂CO₃ (powder). The mixture was extracted with ethylacetate/CH₃OH. The organic layer was separated, dried (over MgSO₄), filtered and the solvent was evaporated until dryness, yielding 2.6g of intermediate **e-9** (87%). In an analogous way, **e-10** was prepared.


A mixture of **e-9** (0.0065 mol) and Raney Nickel (2.6g) in CH₃OH (100ml) was hydrogenated at room temperature for 1 hour under a 3 bar pressure, and then filtered over celite. Celite was washed with CH₃OH. The filtrate was evaporated. The residue (2.2g) was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH/

NH₄OH 85/14/1; 15-40μm). The pure fractions were collected and the solvent was evaporated, yielding 0.85g of intermediate **c-11** (35%). In an analogous way, **c-12** was prepared.


- 5 A mixture of **c-11** (0.000125 mol) and 3-methyl-benzoic acid (0.00025 mol) in CH₂Cl₂ (4 ml) was stirred at room temperature. 1-hydroxybenzotriazole (0.00025 mol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.00025 mol) were added. The reaction was stirred at room temperature for 12 hours. The solution was concentrated and a 10% solution of NaHCO₃ in water (2 ml) and CH₃OH (2 ml) were
- 10 added. The mixture was stirred and refluxed for 4 hours. CH₃OH was then removed under reduced pressure and the resulting solution extracted with CH₂Cl₂. The organic layer was separated, dried (over MgSO₄), filtered and the solvent was evaporated until dryness. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH/NH₄OH 90/10/0.5; 15-40μm). The pure fractions were collected and
- 15 the solvent was evaporated, yielding 0.04g of N-[3-(3-hydroxy-6-methyl-pyridin-2-yl-methyl)-2-(1-methyl-piperidin-4-ylamino)-3H-benzoimidazol-5-yl]-3-methyl-benzamide (60%, compound 65). In an analogous way, N-[1-(3-hydroxy-6-methyl-pyridin-2-ylmethyl)-2-(1-methyl-piperidin-4-ylamino)-1H-benzoimidazol-5-yl]-3-methyl-benzamide (0.028 g or 42% yield, compound 78) was prepared.

20 Example 6

Scheme F



- A mixture of **f-1** (0.0005 mol), 3-methyl-benzaldehyde (0.0006 mol), BH₃CN on solid support (0.0006 mol) and acetic acid (8 drops) in CH₃OH (10ml) was stirred at room
- 25 temperature for 24 hours. The solid support was filtered and washed with CH₃OH. The filtrate was evaporated. The residue (0.53g) was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH/NH₄OH 92/8/0.5 to 89/10/1; 10μm). The pure fractions were collected and the solvent was evaporated. The residue (0.11g) was crystallized from CH₃OH/disopropylether. The precipitate was separated and dried,
- 30 yielding 0.072g of compound **f-2**, i.e. 6-methyl-2-[6-(3-methyl-benzylamino)-2-(1-methyl-piperidin-4-ylamino)-benzoimidazol-1-ylmethyl]-pyridin-3-ol (28%, compound 28, melting point: 240°C).

Example 7Scheme G

5

A mixture of **g-1** (0.0079 mol), **g-2** (0.0095 mol) and triethylamine (0.0118 mol) in dimethylformamide (60ml) was stirred at 80°C for 12 hours. The solvent was evaporated until dryness. The residue was taken up in $\text{CH}_2\text{Cl}_2/\text{H}_2\text{O}$. The organic layer was separated, dried (over MgSO_4), filtered and the solvent was evaporated until dryness. The residue (7g) was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$ 90/10/0.5; 15-40 μm). The pure fractions were collected and the solvent was evaporated, yielding 1.2g of intermediate **g-3** (30%).

A mixture of **g-3** (0.0023 mol) and MnO_2 (2.4g) in CH_2Cl_2 (80ml) was stirred at room temperature for 12 hours, and then filtered over celite. Celite was washed with H_2O . The filtrate was evaporated until dryness. The residue (1.2g) was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$ 95/5/0.1; 35-70 μm).

15

The pure fractions were collected and the solvent was evaporated, yielding 0.8g of intermediate **g-4** (67%).

Diethyl benzyl phosphonate (0.0023 mol) was added to a mixture of NaH (0.0047 mol) in tetrahydrofuran (20ml) at 5°C under N₂ flow. The mixture was stirred at 5°C for 30 minutes. A solution of **g-4** (0.0007 mol) in tetrahydrofuran (10ml) was added drop wise. The mixture was stirred at 5°C for 1 hour, and then stirred at room temperature for 12 hours. H₂O was added. The mixture was extracted with CH₂Cl₂. The organic layer was separated, dried (over MgSO₄), filtered and the solvent was evaporated until dryness. The residue was crystallized from CH₃OH/2-propanone. The precipitate was filtered off and dried, yielding 0.24g of intermediate **g-5** (52%).


A mixture of **g-5** (0.0001 mol) in a 5N solution of HCl in 2-propanol (0.5ml) and 2-propanol (5ml) was stirred at 60°C for 4 hours, and then cooled to room temperature. The precipitate was filtered, washed with 2-propanol/diisopropylether and dried, yielding 0.058g of 2-{2-[1-(2-Amino-ethyl)-piperidin-4-ylamino]-6-styryl-benzoimidazol-1-ylmethyl}-6-methyl-pyridin-3-ol hydrochloride salt (**g-6**; compound 143, 63%, 3.69 HCl + 3.03 H₂O, melting point: > 260°C).

A mixture of **g-5** (0.0002 mol) and Pd/C 10% (0.03g) in CH₃OH (10ml) and tetrahydrofuran (10ml) was hydrogenated at room temperature for 4 hours under a 2 bar pressure, and then filtered over celite. Celite was washed with H₂O. The filtrate was evaporated until dryness, yielding 0.14g of intermediate **g-7** (100%). This product was used directly in the next reaction step.


A mixture of **g-7** (0.0002 mol) in a 5N solution of HCl in 2-propanol (1.4ml) and 2-propanol (15ml) was stirred at 60°C for 4 hours, and then cooled to room temperature. The precipitate was filtered, washed with 2-propanol/diisopropylether and dried, yielding 0.138g of **g-8**, 2-{2-[1-(2-amino-ethyl)-piperidin-4-ylamino]-6-phenethyl-benzoimidazol-1-ylmethyl}-6-methyl-pyridin-3-ol hydrochloride salt (87%, 3.62 HCl + 2.41 H₂O, compound 140, melting point: 181°C).

Example 8

Scheme H



-50-



A mixture of **h-1** (0.0027 mol), ethyl-bromo acetate (0.0032 mol) and triethylamine
 5 (0.004 mol) in dimethylformamide (40ml) was stirred at 50°C for 1 hour, poured into
 ice water and extracted three times with CH₂Cl₂. The organic layer was separated, dried
 (over MgSO₄), filtered and the solvent was evaporated until dryness. The residue was
 taken up in 2-propanone/diisopropylether. The precipitate was filtered, washed with
 10 H₂O and dried, yielding 1g of intermediate **h-2** (82%).

Intermediate **h-3** was prepared analogous to the procedure described for the preparation
 of **g-4**.

CH₃CO₂H (0.2ml) was added at room temperature to a mixture of **h-3** (0.0004 mol),
 15 3-methyl-aniline (0.0005 mol) and NaBH₃CN (0.0005 mol) in CH₃CN (20ml). The
 mixture was stirred at room temperature for 6 hours. CH₃CO₂H (0.2ml) was added. The
 mixture was stirred at room temperature for 12 hours. The solvent was evaporated until
 dryness. The residue was taken up in CH₂Cl₂/K₂CO₃ 10%. The organic layer was
 separated, dried (over MgSO₄), filtered and the solvent was evaporated until dryness,
 20 yielding 0.22g of intermediate **h-4**(100%). This product was used directly in the next
 reaction step.


LiAlH₄ (0.0008 mol) was added to a mixture of **h-4** (0.0004 mol) in tetrahydrofuran
 (20ml) at 5°C under N₂ flow. The mixture was stirred at 5°C for 1 hour, then brought to
 25 room temperature and stirred for 4 hours. A minimum of H₂O and then CH₂Cl₂ were
 added. The organic layer was separated, dried (over MgSO₄), filtered and the solvent
 was evaporated until dryness. The residue (0.22g) was purified by column chromatog-
 raphy over silica gel (eluent: CH₂Cl₂/CH₃OH/NH₄OH 85/15/1; 15-40μm). The pure
 fractions were collected and the solvent was evaporated. The residue (0.1g, 50%) was

crystallized from CH₃OH/CH₃CN/diisopropylether. The precipitate was filtered off and dried, yielding 0.08g of 2-[2-[1-(2-hydroxy-ethyl)-piperidin-4-ylamino]-6-(m-tolyl-amino-methyl)-benzoimidazol-1-ylmethyl]-6-methyl-pyridin-3-ol (40%, compound 145, melting point: 137°C).

5

Example 9

Scheme I



10


A mixture of **i-1** (0.0185 mol) in ethanol (60ml) and H₂SO₄ 36N (5ml) was stirred and refluxed for 24 hours. The solvent was evaporated until dryness. The residue was taken up in CH₂Cl₂. The organic layer was washed with a 10% solution K₂CO₃ in water, dried (over MgSO₄), filtered and the solvent was evaporated until dryness, yielding
5 3.2g of intermediate **i-2** (89%).

Intermediate **i-3** was prepared in an analogous way to the procedure described for intermediate **a-2**. Intermediate **i-4** was prepared in an analogous way to the procedure described for intermediate **a-3**. Intermediate **i-5** was prepared in an analogous way to
10 the procedure described for intermediate **a-5**.

A mixture of **i-5** (0.0048 mol), 2-chloromethyl-6-methyl-3-pyridinol (0.0058 mol) and K₂CO₃ (0.0192 mol) in dimethylformamide (20ml) was stirred at room temperature for 12 hours, poured into ice water and extracted with CH₂Cl₂. The organic layer was
15 separated, dried (over MgSO₄), filtered and the solvent was evaporated until dryness. The residue (3.4g) was purified by column chromatography over silica gel (eluent: toluene/2-propanol/NH₄OH 83/16/1; 15-35µm). Two fractions were collected and the solvent was evaporated, yielding 0.9g of intermediate **i-6** (37%) and 0.78g of
intermediate **i-7** (32%).

Intermediate **i-8** was prepared in an analogous way to the procedure described for intermediate **a-9**. Intermediate **i-9** was prepared in an analogous way to the procedure described for intermediate **a-11**. Intermediate **i-10** (melting point: 221°C) was prepared
20 in an analogous way to the procedure described for intermediate **h-2**. Intermediate **i-11**
25 was prepared in an analogous way to the procedure described for intermediate **h-3**. Intermediate **i-12** (melting point: 143°C) was prepared in an analogous way to the procedure described for intermediate **h-4**.

2-[2-[1-(2-Hydroxy-ethyl)-piperidin-4-ylamino]-4-methyl-6-(m-tolylamino-methyl)-benzoimidazol-1-ylmethyl]-6-methyl-pyridin-3-ol (**i-13**, compound 168, melting point: 123°C) was prepared in an analogous way to the procedure described for compound
30 **h-5**.

Example 10Scheme J

Intermediate **j-2** was prepared in an analogous way to the procedure described for intermediate **h-3**.

- 10 Diethyl cyanomethyl phosphonate (0.0052 mol) was added to a mixture of NaH (0.0105 mol) in tetrahydrofuran (30ml) at 5°C under N₂ flow. The mixture was stirred at 5°C for 30 minutes. A solution of **j-2** (0.0017 mol) in tetrahydrofuran (20ml) was then added. The mixture was stirred at 5°C for 1 hour, and then stirred at room temperature for 12 hours. H₂O was added. The mixture was extracted with CH₂Cl₂. The
- 15 organic layer was separated, dried (over MgSO₄), filtered and the solvent was evaporated until dryness. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH/NH₄OH 95/5/0.1; 70-200μm). The pure fractions were collected and the solvent was evaporated, yielding 0.7g of intermediate **j-3** (84%).
- 20 A mixture of **j-3** (0.0014 mol) and Pd/C 10% (0.25g) in CH₃OH (35ml) was hydrogenated at 40°C for 6 hours under an 8 bar pressure, then cooled to room

temperature and filtered over celite. The filtrate was evaporated until dryness, yielding 0.3g of intermediate **j-4** (52%).


- Intermediate **j-5** was prepared in an analogous way to the procedure described for intermediate **h-2**. Intermediate **j-6** (melting point: 207°C) was prepared in an analogous way to the procedure described for intermediate **i-13**.

A mixture of **j-6** (0.0003 mol) and Raney Nickel (0.2g) in a saturated solution of NH₃ in CH₃OH (25ml) was hydrogenated at room temperature for 1 hour, and then filtered over celite. The filtrate was evaporated until dryness. The residue (0.22g) was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH/NH₄OH 80/20/2; 10µm). The pure fractions were collected and the solvent was evaporated. The residue (0.083g, 49%) was dissolved in ethanol/2-propanone and converted into the hydrochloric acid salt. The precipitate was filtered off and dried. The residue was crystallized from diisopropylether. The precipitate was filtered off and dried, yielding 0.08g of 2-{6-(3-Amino-propyl)-2-[1-(2-hydroxy-ethyl)-piperidin-4-ylamino]-benzoimidazol-1-ylmethyl}-6-methyl-pyridin-3-ol hydrochloride salt (36%, 3.6 HCl, compound 157, melting point: 185°C).

Example 11

20

Scheme K



- A mixture of **k-1** (0.0019 mol), 3-chloro-propylsulfonamide (0.0022 mol) and triethylamine (0.0028 mol) in dimethylformamide (50ml) was stirred at 70°C for 48 hours, poured into ice water, saturated with K₂CO₃ (powder) and extracted with CH₂Cl₂. The organic layer was separated, dried (over MgSO₄), filtered and the solvent was evaporated until dryness. The residue (1.5g) was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH/NH₄OH 85/15/2; 15-40µm). The pure


fractions were collected and the solvent was evaporated, yielding 0.336g of intermediate **k-2** (36%).

5 A mixture of **k-2** (0.0007 mol) and MnO_2 (1g) in CH_2Cl_2 (30ml) was stirred at room temperature for 6 hours, and then filtered over celite. Celite was washed with H_2O . The solvent of the filtrate was evaporated until dryness, yielding 0.33g of intermediate **k-3** (100%). This product was used directly in the next reaction step.


10 $\text{CH}_3\text{CO}_2\text{H}$ (0.2ml) was added at room temperature to a mixture of **k-3** (0.0004 mol), 3,5-dimethyl-aniline (0.0005 mol) and NaBH_3CN (0.0005 mol) in CH_3CN (20ml). The mixture was stirred at room temperature for 30 minutes. $\text{CH}_3\text{CO}_2\text{H}$ (0.2ml) was added. The mixture was stirred at room temperature for 12 hours. The solvent was evaporated until dryness. The residue was taken up in CH_2Cl_2 . The organic layer was washed with a 10% solution of K_2CO_3 in water, dried (over MgSO_4), filtered and the solvent was
15 evaporated until dryness. The residue (0.26g) was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$ 90/10/1; $5\mu\text{m}$). The pure fractions were collected and the solvent was evaporated, yielding 0.09g (32%). This fraction was crystallized from CH_3CN /diisopropylether. The precipitate was filtered off and dried, yielding 0.083g of 2-{4-[6-[(3,5-Dimethyl-phenylamino)-methyl]-1-(3-hydroxy-6-
20 methyl-pyridin-2-ylmethyl)-1H-benzoimidazol-2-ylamino]-piperidin-1-yl}-ethane-sulfonic acid amide (30%, compound 147, melting point: 142°C).

Example 12

Scheme L



-56-




Intermediate **I-2** (melting point: 210°C) was prepared in an analogous way to the procedure described for intermediate **k-2**. Intermediate **I-3** was prepared in an analogous way to the procedure described for intermediate **k-3**. Intermediate **I-4** was prepared in an analogous way to the procedure described for compound **k-4**.

A mixture of **I-4** (0.0003 mol) in a 7N solution of NH_3 in CH_3OH (15ml) was stirred at 80°C for 12 hours. The solvent was evaporated until dryness. The residue (0.21g) was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$ 85/14/1; 10 μm). The pure fractions were collected and the solvent was evaporated. The residue (0.057g, 30%) was crystallized from 2-propanone/ CH_3CN /diisopropylether. The precipitate was filtered off and dried, yielding 0.05g of 2-{4-[6-[(3,5-dimethylphenylamino)-methyl]-1-(3-hydroxy-6-methyl-pyridin-2-ylmethyl)-1H-benzimidazol-2-ylamino]-piperidin-1-yl}-acetamide (26%, compound 148, melting point: 206°C).

Example 13

Scheme M




A mixture of **m-1** (0.0002 mol) in tetrahydrofuran (30ml) was cooled to 5°C under N_2 flow. LiAlH_4 (0.0007 mol) was added. The mixture was stirred at 5°C for 1 hour, and then stirred at room temperature for 1 hour. A minimum of H_2O was added. CH_2Cl_2 was added. The organic layer was separated, dried (over MgSO_4), filtered and the solvent was evaporated until dryness. The residue (0.16g) was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$ 85/15/1; 10 μm). The pure fractions were collected and the solvent was evaporated. The residue (0.073g, 53%) was crystallized from 2-propanone/ CH_3CN /diisopropylether. The precipitate was filtered off and dried, yielding 0.064g of 2-{6-[(3,5-dimethylphenylamino)-methyl]-2-

[1-(2-hydroxy-ethyl)-piperidin-4-ylamino]-benzoimidazol-1-ylmethyl}-6-methyl-pyridin-3-ol (**m2**, 46%, compound 149, melting point: 144°C).

Example 14

5

Scheme N



10

A mixture of **n-1** (0.022 mol) and N-(propylamino)-piperidine (0.0207 mol) was stirred at 140°C for 1 hour, and then taken up in CH_2Cl_2/CH_3OH . The organic layer was washed with 10% K_2CO_3 , dried (over $MgSO_4$), filtered and the solvent was evaporated until dryness. The residue was purified by column chromatography over silica gel (eluent: $CH_2Cl_2/CH_3OH/NH_4OH$ 92/8/1; 70-200 μ m). The pure fractions were collected and the solvent was evaporated, yielding 2.2g of intermediate **n-2** (30%).

15

A mixture of **n-2** (0.0066 mol), **n-3** (0.0073 mol) and K_2CO_3 (0.02 mol) in dimethylformamide (25ml) was stirred at room temperature for 24 hours, poured into H_2O and extracted with CH_2Cl_2 . The organic layer was separated, dried (over $MgSO_4$),


20

filtered and the solvent was evaporated until dryness. The residue was taken up in CH₃CN/diisopropylether. The precipitate was filtered, washed with H₂O and dried, yielding 1.8g of the mixture of intermediates **n-4** and **n-5** (61%).

- 5 LiAlH₄ (0.012 mol) was added portion wise to a mixture of **n-4** (0.002 mol) and **n-5** (0.002 mol) in tetrahydrofuran (60ml) at 5°C under N₂ flow. The mixture was stirred at 5°C for 1 hour, then at room temperature for 12 hours. A minimum of H₂O was added. A solution of CH₂Cl₂/CH₃OH (90/10) was added. The organic layer was separated, dried (over MgSO₄), filtered and the solvent was evaporated until dryness. The residue
- 10 (1.65g) was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH/NH₄OH 80/20/3; 15-40µm). Two fractions were collected and the solvent was evaporated, yielding 0.35g of fraction 1 and 0.049g of fraction 2. Fraction 1 was crystallized from 2-propanone/diisopropylether. The precipitate was filtered off and dried, yielding 0.33g of intermediate **n-6** (19%, melting point: 220°C). Fraction 2 was
- 15 crystallized from 2-propanone/diisopropylether. The precipitate was filtered off and dried, yielding 0.43g of intermediate **n-7** (26%, melting point: 146°C).

- A mixture of **n-6** (0.0006 mol) and MnO₂ (0.5g) in CH₂Cl₂ (30ml) was stirred at room temperature for 12 hours, and then filtered over celite. Celite was washed with H₂O.
- 20 The filtrate was evaporated until dryness, yielding 0.26g of intermediate **n-8** (100%). The compound was used directly in the next reaction step.


- CH₃CO₂H (0.3ml) was added to a mixture of **n-8** (0.0006 mol), 3,5-dimethyl-aniline (0.0007 mol) and NaBH₃CN (0.0007 mol) in CH₃CN (30ml). The mixture was stirred
- 25 at room temperature for 30 minutes. CH₃CO₂H (0.3ml) was added. The mixture was stirred at room temperature for 24 hours. The solvent was evaporated until dryness. The residue was taken up in 2-propanone/HCl 5N/ethanol. The mixture was stirred at 80°C for 12 hours. The solvent was evaporated until dryness. The mixture was extracted with CH₂Cl₂. The organic layer was separated, dried (over MgSO₄), filtered and the solvent
- 30 was evaporated until dryness. The residue (0.39g) was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH/NH₄OH 90/10/0.5; 10µm). The pure fractions were collected and the solvent was evaporated. The residue (0.119g, 59%) was taken up in CH₃CN/diisopropylether. The precipitate was filtered off and dried, yielding 0.17g of 2-[6-[(3,5-dimethyl-phenylamino)-methyl]-2-(3-piperidin-1-yl-propylamino)-benzoimidazol-1-ylmethyl]-6-methyl-pyridin-3-ol (**n-9**, 53%, compound
- 35 170, melting point: 161°C).

Example 15Scheme O


- 5 2-{6-[(3,5-Dimethyl-phenylamino)-methyl]-2-[3-(4-methyl-piperazin-1-yl)-propylamino]-benzoimidazol-1-ylmethyl}-6-methyl-pyridin-3-ol (compound 171, melting point: 150°C) and its intermediates were prepared in an analogous way to the procedures described for preparing compound **n-9**.

Example 16

10

Scheme P

-60-




Intermediate **p-9** (melting point: 212°C) was prepared in an analogous way to the procedure described for compound **n-9**.

5


A mixture of **p-9** (0.0004 mol) in a 3N solution of HCl in water (20ml) and tetrahydrofuran (20ml) was stirred at room temperature for 6 hours, basified with K_2CO_3 (powder) and extracted with CH_2Cl_2 . The organic layer was separated, dried (over $MgSO_4$), filtered and the solvent was evaporated until dryness. The residue (0.25g) was purified by column chromatography over silica gel (eluent: $CH_2Cl_2/CH_3OH/NH_4OH$ 92/8/0.5; 10 μ m). The pure fractions were collected and the solvent was evaporated. The residue (0.17g, 92%) was crystallized from CH_3CN /diisopropylether. The precipitate was filtered off and dried, yielding 0.127g of 3-[6-[(3,5-dimethyl-phenyl-amino)-methyl]-1-(3-hydroxy-6-methyl-pyridin-2-ylmethyl)-1H-benzoimidazol-2-ylamino]-propane-1,2-diol (**p-10**, 69%, compound 172, melting point: 128°C).

15












Example 17**Scheme Q**













- 10 2-{{6-[(3,5-Dimethyl-phenylamino)-methyl]-2-[2-(4-methoxy-phenyl)-ethylamino]-benzoimidazol-1-ylmethyl}}-6-methyl-pyridin-3-ol (compound 187, melting point: 178°C) and its intermediates were prepared in an analogous way to the procedures described for preparing compound n-9.













- 15 The following tables list compounds that were prepared according to any one of the above examples.














Table 1













Comp. No.	R2b	R2a	Activity	Mass (MH ⁺)	Melting point/salt	Synthesis scheme
1	H		9.4	529	201°C	A
2	H		8.7	525	—	B Variant 1
3	H		8.6	485	—	B Variant 1
4	H		8.5	501	229°C	A
5	H		8.5	556	199°C	A
6	H		8.4	535	—	B Variant 1
7	H		8.4	487	—	B Variant 1
8	H		8.4	485	—	B Variant 1
9	H		8.3	471	141°C	B Variant 2
10	H		8.3	536	—	B Variant 1













Comp. No.	R2b	R2a	Activity	Mass (MH ⁺)	Melting point/salt	Synthesis scheme
11	H		8.2	491	—	B Variant 1
12	H		8.2	482	—	B Variant 1
13	H		8.1	481	—	B Variant 1
14	H		8.1	500	—	B Variant 1
15	H		8	485	150°C	B Variant 2
16	H		8	549	176°C	A
17	H		8	499	—	B Variant 1
18	H		7.9	475	130°C	B Variant 2
19	H		7.9	499	—	B Variant 1
20	H		7.9	485	—	B Variant 1
21	H		7.9	499	—	B Variant 1














Comp. No.	R2b	R2a	Activity	Mass (MH ⁺)	Melting point/salt	Synthesis scheme
22	H		7.9	535	—	B Variant 1
23	H		7.9	475	—	B Variant 1
24	H		7.9	545	—	B Variant 1
25	H		7.9	503	—	B Variant 1
26	H		7.9	475	—	B Variant 1
27	H		7.9	571	197°C	A
28	H		7.8	471	240°C	F
29	H		7.8	456	232°C	G
30	H		7.8	457	206	B Variant 2
31	H		7.8	508	—	B Variant 1
32	H		7.8	515	—	B Variant 1
33	H		7.8	487	—	B Variant 1













Comp. No.	R2b	R2a	Activity	Mass (MH ⁺)	Melting point/salt	Synthesis scheme
34	H		7.8	496	—	B Variant 1
35	H		7.7	470		G
36	H		7.7	517	—	B Variant 1
37	H		7.7	471	—	B Variant 1
38	H		7.7	535	—	B Variant 1
39	H		7.7	515		A
40	H		7.6	505		B Variant 1
41	H		7.6	485	—	B Variant 1
42	H		7.6	533	—	B Variant 1
43	H		7.6	510	—	B Variant 1
44	H		7.5	409	—	J
45	H		7.5	497	—	B Variant 1
















Comp. No.	R2b	R2a	Activity	Mass (MH ⁺)	Melting point/salt	Synthesis scheme
46	H		7.5	493	—	B Variant 1
47	H		7.5	499	—	B Variant 1
48	H		7.5	553	—	B Variant 1
49	H		7.4	514	—	B Variant 1
50	H		7.4	493	—	B Variant 1
51	H		7.4	471	—	B Variant 1
52	H		7.4	513	—	B Variant 1
53	H		7.4	514	—	B Variant 1
54	H		7.4	529	—	B Variant 1
55		H	7.3	472	—	E
56	H		7.3	457	225°C	F
57	H		7.3	529	205°C	B Variant 2
58	H		7.3	500	—	B Variant 1















Comp. No.	R2b	R2a	Activity	Mass (MH ⁺)	Melting point/salt	Synthesis scheme
59	H		7.3	542	185°C / HCl salt	A
60	H		7.3	489	—	B Variant 1
61	H		7.3	517	—	B Variant 1
62	H		7.3	485	233°C	A
63	H		7.3	549	—	B Variant 1
64	H		7.3	548	—	B Variant 1
65	H		7.2	485	—	E
66	H		7.2	549	—	B Variant 1
67	H		7.2	505	—	B Variant 1
68	H		7.2	549	—	B Variant 1
69	H		7.2	553	—	B Variant 1
70	H		7.2	525	—	B Variant 1

Comp. No.	R2b	R2a	Activity	Mass (MH ⁺)	Melting point/salt	Synthesis scheme
71	H		7.2	525	—	B Variant 1
72	H		7.1	529	—	B Variant 1
73	H		7.1	505	—	B Variant 1
74	H		7.1	563	—	B Variant 1
75	H		7	485	—	B Variant 1
76	H		7	499	—	B Variant 1
77	H		7	559	—	B Variant 1
78		H	6.9	485	—	E
79	H		6.9	432	—	A
80	H		6.9	547	249°C	A
81	H		6.9	561	184°C	A
82	H		6.9	513	—	B variant 1

Comp. No.	R2b	R2a	Activity	Mass (MH ⁺)	Melting point/salt	Synthesis scheme
83	H		6.9	569	—	B Variant 1
84	H		6.8	470	224°C	G
85	H		6.8	515	189°C	A
86	H		6.8	458	144°C	
87	H		6.8	563	—	B Variant 1
88	H		6.8	519	—	B Variant 1
89	H		6.8	569	—	B Variant 1
90		H	6.7	471	—	E
91	H		6.7	517	—	B Variant 1
92	H		6.6	382	165°C	A
93	H		6.6	488	202°C	C
94	H		6.6	585	—	B Variant 1
95	H		6.6	547	—	A

Comp. No.	R2b	R2a	Activity	Mass (MH ⁺)	Melting point/salt	Synthesis scheme
96	H		6.6	499	—	B Variant 1
97	H		6.6	495	171°C	A
98	H		6.5	451	224°C	A
99	H		6.4	454	262°C	G
100		H	6.4	454	> 260°C	G
101	H	H	6.4	352	> 260°C	G
102	H		6.4	497	—	A
103	H		6.3	468	244°C	G
104		H	6.2	468	261°C	G
105	H		6.2	499	—	A
106	H		6.1	499	—	E
107	H		6.1	485	—	E
108	H		6	486	—	E

Comp. No.	R2b	R2a	Activity	Mass (MH ⁺)	Melting point/salt	Synthesis scheme
109	H		6	481	—	A
110	H		5.8	485	102°C	B Variant 2
111	H		5.6	453	—	A
112	H		5.5	471	169°C	D
113		H	5.5	486	—	E
114	H		5.2	472	—	E
115		H	5.1	403	188°C	J
116	H		5.1	485	172°C	D
117	H		5.1	494	161°C	B Variant 2
118		H	4.9	475	250°C	D
119	H		4.8	475	155°C	D
120		H	4.6	499	200°C	D
121		H	< 4	424	243°C	D
122		H	< 4	382	> 260°C	A
123		H	4.7	456	159°C	G

Comp. No.	R2b	R2a	Activity	Mass (MH ⁺)	Melting point/salt	Synthesis scheme
124		H	5.1	470	196°C	G
125		H	< 4	468	229°C	G
126	H		< 4	488	248°C	G
127		H	< 4	488	225°C	G
128	H		< 4	468	250°C	G
129		H	< 4	470	222°C	G
130		H	4.6	471	156°C	B Variant 2
131		H	< 4	457	199°C	B Variant 2
132		H	< 4	488	208°C	C
133		H	< 4	409	245°C	J
134	H		< 4	486	146°C	D
135		H	< 4	486	230°C	D
136	H		< 4	499	194°C	D
137	H		< 4	487	—	A



Comp. No.	R2b	R2a	Activity	Mass (MH ⁺)	Melting point/salt	Synthesis scheme
138	H		9.6	570	180°C	A

Table 2a






Comp. No.	R	Activity	Mass (MH ⁺)	Melting point/salt	Synthesis scheme
139		5.1	410	164°C	D
140		< 4	500	238°C	A

Table 2b






Comp. No.	R	Activity	Mass (MH ⁺)	Melting point/salt	Synthesis scheme
141		C	410	262°C	D
142		C	500	179°C	A


Table 3



Comp. No.	R	R ^{2b}	R ^{2a}	Activity	Mass (MH ⁺)	Melting point/salt	Synthesis scheme
143	-CH ₂ -NH ₂	H		9	485	181°C / HCl	G
144	-CH ₂ -NH ₂	H		9	499	188°C / HCl	G
145	-CH ₂ -NH ₂		H	9	483	> 260°C / HCl	G
146	-CH ₂ -NH ₂	H		8.9	483	> 260°C / HCl	G
147	-CH ₂ -NH ₂		H	8.8	497	210°C / HCl	G
148	-CH ₂ -OH	H		8.8	501	137°C	H
149	-CH ₂ -OH	H		8.8	515	133°C	H
150		H		8.6	592	142°C	K
151		H		8.6	542	206°C	L
152	-(CH ₂) ₂ -OH	H		8.6	529	144°C	M
153	-CH ₂ -NH ₂	H		8.5	513	211°C / HCl	G
154		H		8.5	578	193°C	K
155		H		8.5	528	145°C	L


Comp. No.	R	R ^{2b}	R ^{2a}	Activity	Mass (MH ⁺)	Melting point/salt	Synthesis scheme
156	HO-CH ₂ -CH ₂ -CH ₂ -	H		8.4	515	—	M
157	H ₂ N-CH ₂ -		H	7.9	485	183°C / HCl	G
158	H ₂ N-CH ₂ -		H	7.9	499	179°C / HCl	G
159	HO-CH ₂ -	H		7.9	503	158°C	H
160	HO-CH ₂ -	H		7.9	439	185°C / HCl	J
161	HO-CH ₂ -	H		7.8	435	207°C	J
162	HO-CH ₂ -	H		7.5	486	201°C	G
163		H		7.2	513	202°C	G
164	phenyl	H		7.2	561	186°C	H
165	phenyl	H		7.1	601	149°C	H
166			H	6.8	511	—	G
167	phenyl	H		6.8	547	198°C	H
168	HO-CH ₂ -		H	6.6	486	216°C	G
169	HO-CH ₂ -	H		6.6	484	240°C	G
170			H	5.7	513	160°C	G

Table 4


























Comp. No.	R ^{2a}	Activity	Mass (MH ⁺)	Melting point/salt	Synthesis scheme
171		8.5	515	123°C	I
172		8.5	529	136°C	I

Table 5



Comp. No.	R	R ^{2b}	R ^{2a}	Activity	Mass (MH ⁺)	Melting point/salt	Synthesis scheme
173		H		8.8	513	161°C	N
174		H		8.4	528	150°C	O
175		H		7.5	462	128°C	P
176			H	7.2	513	182°C	N
177		H	-CH ₂ -OH	7.1	410	220°C	N

Comp. No.	R	R ^{2b}	R ^{2a}	Activity	Mass (MH ⁺)	Melting point/salt	Synthesis scheme
178		H	-CH ₂ -OH	6.5	425	230°C	O
179			H	6.2	528	193°C	O
180			H	5.6	462	215°C	P
181		H	-CH ₂ -OH	5.1	419	194°C	Q
182		-CH ₂ -OH	H	5	410	146°C	N
183		-CH ₂ -OH	H	5	425	154°C	O
184		H		< 4	502	212°C	P
185		-CH ₂ -OH	H	< 4	389	230°C	Q
186			H	< 4	492	175°C	Q
187		H		< 4	492	190°C	Q
188		H	-CH ₂ -OH	4.4	389	185°C	Q
189		-CH ₂ -OH	H	< 4	419	185°C	Q
190		H		< 4	522	178°C	Q
191			H	< 4	522	196°C	Q

Comp. No.	R	R ^{2b}	R ^{2a}	Activity	Mass (MH ⁺)	Melting point/salt	Synthesis scheme
192		H		< 4	532	211°C	Q

Example 18: *In vitro* screening for activity against Respiratory Syncytial Virus.

The percent protection against cytopathology caused by viruses (antiviral activity or EC₅₀) achieved by tested compounds and their cytotoxicity (CC₅₀) are both calculated from dose-response curves. The selectivity of the antiviral effect is represented by the selectivity index (SI), calculated by dividing the CC₅₀ (cytotoxic dose for 50% of the cells) by the EC₅₀ (antiviral activity for 50 % of the cells). The tables in the above experimental part list the category to which each of the prepared compounds belong : Compounds belonging to activity category "A" have an pEC₅₀ (-log of EC₅₀ when expressed in molar units) equal to or more than 7. Compounds belonging to activity category "B" have a pEC₅₀ value between 6 and 7. Compounds belonging to activity category "C" have a pEC₅₀ value equal to or below 6.


Automated tetrazolium-based colorimetric assays were used for determination of EC₅₀ and CC₅₀ of test compounds. Flat-bottom, 96-well plastic microtiter trays were filled with 180 µl of Eagle's Basal Medium, supplemented with 5 % FCS (0% for FLU) and 20 mM Hepes buffer. Subsequently, stock solutions (7.8 x final test concentration) of compounds were added in 45 µl volumes to a series of triplicate wells so as to allow simultaneous evaluation of their effects on virus- and mock-infected cells. Five five-fold dilutions were made directly in the microtiter trays using a robot system. Untreated virus controls, and HeLa cell controls were included in each test. Approximately 100 TCID₅₀ of Respiratory Syncytial Virus was added to two of the three rows in a volume of 50 µl. The same volume of medium was added to the third row to measure the cytotoxicity of the compounds at the same concentrations as those used to measure the antiviral activity. After two hours of incubation, a suspension (4 x 10⁵ cells/ml) of HeLa cells was added to all wells in a volume of 50µl. The cultures were incubated at 37°C in a 5% CO₂ atmosphere. Seven days after infection the cytotoxicity and the antiviral activity was examined spectrophotometrically. To each well of the microtiter tray, 25 µl of a solution of MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) was added. The trays were further incubated at 37°C for 2 hours, after which the medium was removed from each cup. Solubilization of the formazan crystals was achieved by adding 100 µl 2-propanol. Complete dissolution of the formazan crystals were obtained after the trays have been placed on a plate shaker for 10 min. Finally, the

-79-

absorbances were read in an eight-channel computer-controlled photometer (Multiskan MCC, Flow Laboratories) at two wavelengths (540 and 690 nm). The absorbance measured at 690 nm was automatically subtracted from the absorbance at 540 nm, so as to eliminate the effects of non-specific absorption.

Claims

1. A compound having the formula




a prodrug, *N*-oxide, addition salt, quaternary amine, metal complex or stereochemically isomeric form thereof wherein


Q is Ar², R^{6a}, pyrrolidinyl substituted with R⁶, piperidinyl substituted with R⁶ or homopiperidinyl substituted with R⁶;

G is a direct bond or C₁₋₁₀alkanediyl optionally substituted with one or more substituents individually selected from the group consisting of hydroxy, C₁₋₆alkyloxy, Ar¹C₁₋₆alkyloxy, C₁₋₆alkylthio, Ar¹C₁₋₆alkylthio, HO(-CH₂-CH₂-O)_n, C₁₋₆alkyloxy(-CH₂-CH₂-O)_n and Ar¹C₁₋₆alkyloxy(-CH₂-CH₂-O)_n;


R¹ is Ar¹ or a monocyclic or bicyclic heterocycle being selected from piperidinyl, piperazinyl, pyridyl, pyrazinyl, pyridazinyl, pyrimidinyl, furanyl, tetrahydrofuranyl, thienyl, pyrrolyl, thiazolyl, oxazolyl, imidazolyl, isothiazolyl, pyrazolyl, isoxazolyl, oxadiazolyl, quinolinyl, quinoxalinyl, benzofuranyl, benzothienyl, benzimidazolyl, benzoxazolyl, benzthiazolyl, pyridopyridyl, naphthiridinyl, 1*H*-imidazo[4,5-*b*]pyridinyl, 3*H*-imidazo[4,5-*b*]pyridinyl, imidazo[1,2-*a*]pyridinyl, 2,3-dihydro-1,4-dioxino[2,3-*b*]pyridyl or a radical of formula




(c-1)




(c-2)




(c-3)




(c-4)




(c-5)



(c-6)



(c-7)



(c-8)

;


- wherein each of said monocyclic or bicyclic heterocycles may optionally be substituted with 1 or where possible more, such as 2, 3, 4 or 5, substituents individually selected from the group of substituents consisting of halo, hydroxy, amino, cyano, carboxyl, C₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkylthio, C₁₋₆alkyloxyC₁₋₆alkyl, Ar¹, Ar¹C₁₋₆alkyl, Ar¹C₁₋₆alkyloxy, hydroxyC₁₋₆alkyl, mono-or di(C₁₋₆alkyl)amino, mono-or di(C₁₋₆alkyl)aminoC₁₋₆alkyl, polyhaloC₁₋₆alkyl, C₁₋₆alkylcarbonylamino, C₁₋₆alkyl-SO₂-NR^{4a}-, Ar¹-SO₂-NR^{4a}-, C₁₋₆alkyloxycarbonyl, -C(=O)-NR^{4a}R^{4b}, HO(-CH₂-CH₂-O)_n-, halo(-CH₂-CH₂-O)_n-, C₁₋₆alkyloxy(-CH₂-CH₂-O)_n-, Ar¹C₁₋₆alkyloxy(-CH₂-CH₂-O)_n- and mono-and di(C₁₋₆alkyl)amino(-CH₂-CH₂-O)_n-;
- one of R^{2a} and R^{2b} is cyanoC₁₋₆alkyl, cyanoC₂₋₆alkenyl, Ar³C₁₋₆alkyl, (Ar³)(OH)C₁₋₆alkyl, Het-C₁₋₆alkyl, N(R^{8a}R^{8b})C₁₋₆alkyl, Ar³C₂₋₆alkenyl, Het-C₂₋₆alkenyl, Ar³aminoC₁₋₆alkyl, Het-aminoC₁₋₆alkyl, Het-C₁₋₆alkylamino-C₁₋₆alkyl, Ar³thioC₁₋₆alkyl, Het-thioC₁₋₆alkyl, Ar³sulfonylC₁₋₆alkyl, Het-sulfonyl-C₁₋₆alkyl, Ar³aminocarbonyl, Het-aminocarbonyl, Ar³(CH₂)_naminocarbonyl, Het-(CH₂)_naminocarbonyl, Ar³carbonylamino, Het-carbonylamino, Ar³(CH₂)_ncarbonylamino, Het-(CH₂)_ncarbonylamino, Ar³(CH₂)_namino; and the other one of R^{2a} and R^{2b} is hydrogen;
- in case R^{2a} is hydrogen, then R³ is hydrogen;
- in case R^{2b} is hydrogen, then R³ is hydrogen or C₁₋₆alkyl;
- R^{4a} and R^{4b} can be the same or can be different relative to one another, and are each independently hydrogen or C₁₋₆alkyl; or
- R^{4a} and R^{4b} taken together may form a bivalent radical of formula -(CH₂)_s- wherein s is 4 or 5;
- R⁵ is hydrogen or C₁₋₆alkyl;
- R⁶ is hydrogen or C₁₋₆alkyl optionally substituted with one or more substituents each independently selected from the group consisting of trifluoromethyl, NR^{7a}R^{7b}, C₃₋₇cycloalkyl, Ar², hydroxy, C₁₋₄alkoxy, C₁₋₄alkylthio, Ar²-oxy-, Ar²-thio-, Ar²(CH₂)_noxy, Ar²(CH₂)_nthio, hydroxycarbonyl, aminocarbonyl, C₁₋₄alkyl-carbonyl, Ar²carbonyl, C₁₋₄alkoxycarbonyl, Ar²(CH₂)_ncarbonyl, amino-carbonyloxy, C₁₋₄alkylcarbonyloxy, Ar²carbonyloxy, Ar²(CH₂)_ncarbonyloxy, C₁₋₄alkoxycarbonyl(CH₂)_noxy, mono- or di(C₁₋₄alkyl)aminocarbonyl, mono- or di(C₁₋₄alkyl)aminocarbonyloxy, aminosulfonyl, mono- or di(C₁₋₄alkyl)amino-sulfonyl or a heterocycle selected from the group consisting of pyrrolidinyl, pyrrolyl, dihydropyrrolyl, imidazolyl, triazolyl, piperidinyl, homopiperidinyl, piperazinyl, pyridyl and tetrahydropyridyl, wherein each of said heterocycle may optionally be substituted with oxo or C₁₋₆alkyl;
- R^{6a} is C₁₋₆alkyl substituted with one or more substituents each independently selected from the group consisting of trifluoromethyl, NR^{7a}R^{7b}, C₃₋₇cycloalkyl, Ar²,

- hydroxy, C₁₋₄alkoxy, C₁₋₄alkylthio, Ar²-oxy-, Ar²-thio-, Ar²(CH₂)_noxy, Ar²(CH₂)_nthio, hydroxycarbonyl, aminocarbonyl, C₁₋₄alkylcarbonyl, Ar²carbonyl, C₁₋₄alkoxycarbonyl, Ar²(CH₂)_ncarbonyl, aminocarbonyloxy, C₁₋₄alkylcarbonyloxy, Ar²carbonyloxy, Ar²(CH₂)_ncarbonyloxy, C₁₋₄alkoxycarbonyl(CH₂)_noxy,
- 5 mono- or di(C₁₋₄alkyl)aminocarbonyl, mono- or di(C₁₋₄alkyl)aminocarbonyloxy, aminosulfonyl, mono- or di(C₁₋₄alkyl)aminosulfonyl or a heterocycle selected from the group consisting of pyrrolidinyl, pyrrolyl, dihydropyrrolyl, imidazolyl, triazolyl, piperidinyl, homopiperidinyl, piperazinyl, pyridyl and tetrahydro-
- 10 pyridyl, wherein each of said heterocycle may optionally be substituted with oxo or C₁₋₆alkyl;
- R^{7a} is hydrogen, C₁₋₆alkyl, formyl or C₁₋₆alkylcarbonyl;
- R^{7b} is hydrogen, C₁₋₆alkyl, formyl or C₁₋₆alkylcarbonyl;
- R^{8a} is Ar³, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, C₁₋₆alkoxyC₁₋₆alkyl, cyanoC₁₋₆alkyl, aminoC₁₋₆alkyl, mono- or di(C₁₋₆alkyl)aminoC₁₋₆alkyl, Ar³C₁₋₆alkyl, Het-C₁₋₆alkyl,
- 15 aminocarbonyl-C₁₋₆alkyl, carboxyl-C₁₋₆alkyl;
- R^{8b} is Ar³, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, C₁₋₆alkoxyC₁₋₆alkyl, cyanoC₁₋₆alkyl, aminoC₁₋₆alkyl, mono- or di(C₁₋₆alkyl)aminoC₁₋₆alkyl, Ar³C₁₋₆alkyl, Het-C₁₋₆alkyl;
- each n independently is 1, 2, 3 or 4;
- each m independently is 1 or 2;
- 20 each p independently is 1 or 2;
- Ar¹ is phenyl or phenyl substituted with 1 or more, such as 2, 3 or 4, substituents selected from halo, hydroxy, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, polyhaloC₁₋₆alkyl, and C₁₋₆alkyloxy;
- Ar² is phenyl or phenyl substituted with 1 or more, such as 2, 3 or 4, substituents
- 25 selected from the group consisting of halo, hydroxy, amino, cyano, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, polyhaloC₁₋₆alkyl, aminoC₁₋₆alkyl, C₁₋₆alkyloxy, aminosulfonyl, aminocarbonyl, hydroxycarbonyl, C₁₋₄alkylcarbonyl, mono- or di(C₁₋₄alkyl)amino, mono- or di(C₁₋₄alkyl)aminocarbonyl, mono- or di(C₁₋₄alkyl)aminosulfonyl, mono- or di(C₁₋₄alkyl)aminoC₁₋₆alkyl and
- 30 C₁₋₄alkoxycarbonyl;
- Ar³ is phenyl, naphthalenyl, 1,2,3,4-tetrahydro-naphthalenyl or indanyl, wherein said phenyl, naphthyl, 1,2,3,4-tetrahydro-naphthalenyl or indanyl may optionally and each individually be substituted with one or more, such as 2, 3 or 4, substituents selected from the group consisting of halo, hydroxy, mercapto, amino, cyano,
- 35 C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, Ar¹, hydroxyC₁₋₆alkyl, polyhaloC₁₋₆alkyl, aminoC₁₋₆alkyl, cyanoC₁₋₆alkyl, aminocarbonyl, C₁₋₆alkyloxy, C₁₋₆alkylthio, Ar¹-oxy, Ar¹-thio, Ar¹-amino, aminosulfonyl, aminocarbonyl-C₁₋₆alkyl, hydroxycarbonyl-C₁₋₆alkyl, hydroxycarbonyl, C₁₋₄alkylcarbonyl, mono- or

di(C₁₋₄alkyl)amino, mono- or di(C₁₋₄alkyl)aminocarbonyl, mono- or di(C₁₋₄alkyl)aminosulfonyl, mono- or di(C₁₋₄alkyl)aminoC₁₋₆alkyl, C₁₋₄alkylcarbonylamino and C₁₋₄alkoxycarbonyl;

Het is a heterocycle being selected from tetrahydrofuranyl, tetrahydrothienyl, dioxanyl, dioxolanyl, pyrrolidinyl, pyrrolidinonyl, furanyl, thienyl, pyrrolyl, thiazolyl, oxazolyl, imidazolyl, isothiazolyl, pyrazolyl, isoxazolyl, oxadiazolyl, thiadiazolyl, piperidinyl, homopiperidinyl, piperazinyl, morpholinyl, pyridyl, pyrazinyl, pyridazinyl, pyrimidinyl, tetrahydroquinolinyl, quinolinyl, isoquinolinyl, benzodioxanyl, benzodioxolyl, indolinyl, indolyl, each of said heterocycle may optionally be substituted with oxo, amino, Ar¹, C₁₋₄alkyl, aminoC₁₋₄alkyl, hydroxyC₁₋₆alkyl, Ar¹C₁₋₄alkyl, mono- or di(C₁₋₆alkyl)aminoC₁₋₆alkyl, mono- or di(C₁₋₆alkyl)amino, or with two C₁₋₄alkyl radicals.

2. A compound according to claim 1 wherein the compound has the formula (I-a-1):




wherein Q, R⁵, G and R¹ are as claimed in claim 1; and

Alk is C₁₋₆alkanediyl;

R^{8c} has the same meanings of R^{8a}, as claimed in claim 1, and also may be hydrogen;

R⁹, R¹⁰, R¹¹ independently from one another have the same meanings as the substituents on Ar³ as claimed in claim 1.

3. A compound according to claim 1 wherein the compound has the formula (I-b-1):




wherein Q, R⁵, G and R¹ are as claimed in claim 1; and

Alk is C₁₋₆alkanediyl;

R^{8c} has the same meanings of R^{8a}, as claimed in claim 1, and also may be hydrogen;

R^9 , R^{10} , R^{11} independently from one another have the same meanings as the substituents on Ar^3 as claimed in claim 1.


4. A compound according to claim 1 wherein the compound has the formula (I-c):



5

wherein t , G , R^1 , R^{2a} , R^{2b} , R^3 , R^5 and R^6 are as claimed in claim 1.

5. A compound according to claim 1 wherein the compound has the formula (I-d-2):



10

wherein t , R^5 , R^6 , G and R^1 are as claimed in claim 1; and


Alk is C_{1-6} alkanediyl;

R^{8c} has the same meanings of R^{8a} , as claimed in claim 1, and also may be hydrogen;

R^9 , R^{10} , R^{11} independently from one another have the same meanings as the substituents on Ar^3 as claimed in claim 1.

15

6. A compound according to claim 1 wherein the compound has the formula (I-e-2):



20

wherein t , R^5 , R^6 , G and R^1 are as claimed in claim 1; and

Alk is C_{1-6} alkanediyl;

R^{8c} has the same meanings of R^{8a} , as claimed in claim 1, and also may be hydrogen;

R^9 , R^{10} , R^{11} independently from one another have the same meanings as the substituents on Ar^3 as claimed in claim 1.

7. A compound according to any of claims 4 to 6 wherein t is 2.
8. A compound according to any of claims 1 - 7, wherein G is C₁₋₁₀alkanediyl.
9. A compound according to in any of claims 1 - 7, wherein G is methylene.
10. A compound according to any of claims 1 - 9, wherein R¹ is pyridyl optionally substituted with 1 or 2 substituents independently selected from the group consisting of halo, hydroxy, amino, cyano, carboxyl, C₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkylthio, C₁₋₆alkyloxyC₁₋₆alkyl, Ar¹, Ar¹C₁₋₆alkyl, Ar¹C₁₋₆alkyloxy, hydroxyC₁₋₆alkyl, mono-or di(C₁₋₆alkyl)amino, mono-or di(C₁₋₆alkyl)amino-C₁₋₆alkyl, polyhaloC₁₋₆alkyl, C₁₋₆alkylcarbonylamino, C₁₋₆alkyl-SO₂-NR^{4a}-, Ar¹-SO₂-NR^{4a}-, C₁₋₆alkyloxycarbonyl, -C(=O)-NR^{4a}R^{4b}, HO(-CH₂-CH₂-O)_n-, halo(-CH₂-CH₂-O)_n-, C₁₋₆alkyloxy(-CH₂-CH₂-O)_n-, Ar¹C₁₋₆alkyloxy(-CH₂-CH₂-O)_n- and mono-or di(C₁₋₆alkyl)amino(-CH₂-CH₂-O)_n-.
11. A compound according to any of claims 1 - 9, wherein R¹ is pyridyl substituted with 1 or 2 substituents independently selected from the group consisting of hydroxy and C₁₋₆alkyl.
12. A compound according to any of claims 1 - 9, wherein R¹ is Ar¹, quinolinyl, benzimidazolyl, a radical of formula
- (c-4)
- or pyrazinyl; wherein each of the radicals Ar¹, quinolinyl, benzimidazolyl, (c-4), or pyrazinyl may optionally be substituted with the substituents of said radicals as claimed in claim 1.
13. A compound according to any of claims 1 - 9, wherein R¹ is phenyl optionally substituted with one, two or three radicals selected from the group consisting of halo, hydroxy, C₁₋₆alkyl, C₁₋₆alkyloxy; quinolinyl; a radical (c-4) wherein m is 2, optionally substituted with up to two radicals selected from C₁₋₆alkyl; benzimidazolyl optionally substituted with C₁₋₆alkyl; pyrazinyl optionally substituted with up to three radicals selected from C₁₋₆alkyl.


14. A compound according to any of claims 1 - 13, wherein one of R^{2a} and R^{3a} is selected from cyanoC₁₋₆alkyl, cyanoC₂₋₆alkenyl, Ar³C₁₋₆alkyl, (Ar³)(OH)C₁₋₆alkyl, Het-C₁₋₆alkyl, N(R^{8a}R^{8b})C₁₋₆alkyl, Ar³C₂₋₆alkenyl, Het-C₂₋₆alkenyl, Ar³aminoC₁₋₆alkyl, Het-aminoC₁₋₆alkyl, Het-C₁₋₆alkylaminoC₁₋₆alkyl, Ar³thioC₁₋₆alkyl, Het-thioC₁₋₆alkyl, Ar³sulfonylC₁₋₆alkyl, Het-sulfonylC₁₋₆alkyl, Ar³aminocarbonyl, Het-aminocarbonyl, Ar³(CH₂)_naminocarbonyl, Het-(CH₂)_naminocarbonyl, Ar³carbonylamino, Ar³(CH₂)_namino; and the other one of R^{2a} and R^{2b} is hydrogen.
15. A compound according to any of claims 1 - 13, wherein one of R^{2a} and R^{3a} is selected from cyanoC₁₋₆alkyl, Ar³C₁₋₆alkyl, Het-C₁₋₆alkyl, N(R^{8a}R^{8b})C₁₋₆alkyl, Ar³C₂₋₆alkenyl, Ar³aminoC₁₋₆alkyl, Het-aminoC₁₋₆alkyl, Het-C₁₋₆alkylaminoC₁₋₆alkyl, Ar³thioC₁₋₆alkyl, Ar³aminocarbonyl, Het-aminocarbonyl, Ar³(CH₂)_naminocarbonyl, Het-(CH₂)_naminocarbonyl; and the other one of R^{2a} and R^{2b} is hydrogen.
16. A compound according to any of claims 1 - 13, wherein one of R^{2a} and R^{3a} is selected from N(R^{8a}R^{8b})C₁₋₆alkyl, Ar³aminoC₁₋₆alkyl; and the other one of R^{2a} and R^{2b} is hydrogen.
17. A compound according to any of claims 14 - 16, wherein
in case R^{2a} is hydrogen then R^3 is hydrogen;
in case R^{2b} is hydrogen then R^3 is hydrogen or C₁₋₆alkyl.
18. A compound according to any of claims 1 - 17, wherein R^5 is hydrogen.
19. A compound according to any of claims 1 - 18, wherein Q is R^{6a} , wherein R^{6a} is C₁₋₆alkyl substituted with one or with two substituents each independently selected from the group consisting of trifluoromethyl, NR^{7a}R^{7b}, Ar², hydroxy, C₁₋₄alkoxy, Ar²(CH₂)_noxy, hydroxycarbonyl, aminocarbonyl, C₁₋₄alkylcarbonyl, C₁₋₄alkoxycarbonyl, Ar²(CH₂)_ncarbonyl, aminocarbonyloxy, C₁₋₄alkylcarbonyloxy, Ar²carbonyloxy, mono- or di(C₁₋₄alkyl)aminocarbonyl, aminosulfonyl, mono- or di(C₁₋₄alkyl)aminosulfonyl or a heterocycle selected from the group consisting of pyrrolidinyl, imidazolyl, piperidinyl, homopiperidinyl, piperazinyl, dioxolanyl, dioxanyl and pyridyl, wherein each of said heterocycle may optionally be substituted with one or two radicals selected from oxo and C₁₋₆alkyl;

20. A compound according to any of claims 1 - 18, wherein Q is R^{6a}, wherein R^{6a} is C₁₋₆alkyl substituted with Ar² or hydroxy, or C₁₋₆alkyl substituted with two hydroxy radicals, or C₁₋₆alkyl substituted with diC₁₋₆alkyl-dioxolanyl, pyrrolidinyl, piperidinyl, piperazinyl, 4-C₁₋₆alkyl-piperazinyl.
21. A compound according to any of claims 1 - 18, wherein Q is pyrrolidinyl substituted with R⁶, piperidinyl substituted with R⁶ or homopiperidinyl substituted with R⁶; wherein R⁶ is hydrogen or C₁₋₆alkyl optionally substituted with one or with two substituents, each independently selected from the group consisting of trifluoromethyl, NR^{7a}R^{7b}, Ar², hydroxy, C₁₋₄alkoxy, Ar²(CH₂)_noxy, hydroxycarbonyl, aminocarbonyl, C₁₋₄alkylcarbonyl, C₁₋₄alkoxycarbonyl, Ar²(CH₂)_ncarbonyl, aminocarbonyloxy, C₁₋₄alkylcarbonyloxy, Ar²carbonyloxy, mono- or di(C₁₋₄alkyl)aminocarbonyl, aminosulfonyl, mono- or di(C₁₋₄alkyl)-aminosulfonyl or a heterocycle selected from the group consisting of pyrrolidinyl, imidazolyl, piperidinyl, homopiperidinyl, piperazinyl, dioxolanyl, dioxanyl and pyridyl, wherein each of said heterocycle may optionally be substituted with one or two radicals selected from oxo and C₁₋₆alkyl.
22. A compound according to any of claims 1 - 18, wherein Q is pyrrolidinyl substituted with R⁶, piperidinyl substituted with R⁶ or homopiperidinyl substituted with R⁶; wherein R⁶ is hydrogen or C₁₋₆alkyl optionally substituted with NR^{7a}R^{7b}, Ar², hydroxy, hydroxycarbonyl, aminocarbonyl, aminosulfonyl or C₁₋₆alkyl substituted with two hydroxy radicals, or C₁₋₆alkyl substituted with a heterocycle selected from dioxolanyl, pyrrolidinyl, piperidinyl, homopiperidinyl, piperazinyl, wherein each of said heterocycle may optionally be substituted with oxo or with one or two C₁₋₆alkyl radicals.
23. A compound according to any of claims 1 - 18, wherein Q is pyrrolidinyl substituted with R⁶, piperidinyl substituted with R⁶ or homopiperidinyl substituted with R⁶; wherein R⁶ is hydrogen or C₁₋₆alkyl substituted with Ar² or C₁₋₆alkyl substituted with piperidinyl or with piperazinyl.
24. A compound according to any of claims 21 - 23, wherein Q is piperidinyl substituted with R⁶.
25. A compound according to any of claims 1 - 24, wherein R^{8a} is Ar³, C₁₋₆alkyl, hydroxyc₁₋₆alkyl, C₁₋₆alkoxyc₁₋₆alkyl, cyanoC₁₋₆alkyl, aminoC₁₋₆alkyl, mono-or


di(C₁₋₆alkyl)aminoC₁₋₆alkyl, Ar³C₁₋₆alkyl, Het-C₁₋₆alkyl, aminocarbonyl-C₁₋₆alkyl, carboxyl-C₁₋₆alkyl; and R^{8b} is Ar³.

26. A compound according to any of claims 1 - 24, wherein R^{8a} is C₁₋₆alkyl, hydroxyC₁₋₆alkyl, Ar³C₁₋₆alkyl, Het-C₁₋₆alkyl, aminocarbonyl-C₁₋₆alkyl; and R^{8b} is C₁₋₆alkyl, hydroxyC₁₋₆alkyl, Ar³C₁₋₆alkyl, Het-C₁₋₆alkyl.
27. A compound according to any of claims 1 - 26, wherein Ar³ is phenyl optionally substituted with one, two or three substituents selected from halo, hydroxy, mercapto, amino, cyano, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, Ar¹, hydroxy-C₁₋₆alkyl, CF₃, aminoC₁₋₆alkyl, cyanoC₁₋₆alkyl, aminocarbonyl, C₁₋₆alkyloxy, C₁₋₆alkylthio, Ar¹-oxy, Ar¹-thio, Ar¹-amino, aminosulfonyl, aminocarbonyl-C₁₋₆alkyl, hydroxycarbonyl-C₁₋₆alkyl, hydroxycarbonyl, C₁₋₄alkylcarbonyl, C₁₋₄alkylcarbonylamino or C₁₋₄alkoxycarbonyl.
28. A compound according to any of claims 1 - 27, wherein Ar³ is phenyl substituted with one, two or three substituents selected from halo, C₁₋₆alkyl or hydroxyC₁₋₆alkyl.
29. A compound as claimed in any one of claims 1 to 28 for use as a medicine.
30. A pharmaceutical composition comprising a pharmaceutically acceptable carrier, and as active ingredient a therapeutically effective amount of a compound as claimed in any one of claims 1 to 23.
31. A process for preparing a pharmaceutical composition as claimed in claim 25, said process comprising intimately mixing a pharmaceutically acceptable carrier with a therapeutically effective amount of a compound as claimed in any one of claims 1 to 23.
32. The use of a compound as claimed in any of claims 1 to 23 for the manufacture of a medicament for inhibiting RSV replication.
33. A process for preparing a compound as claimed in any of claims 1 to 23, said process comprising
 - (a) reacting an intermediate of formula (II) with a reagent (III) as in the following reaction scheme:

-89-




(b) reacting an intermediate of formula (IV) with a reagent (V) thus obtaining a compound of formula (I-c-1) as in the following reaction scheme:




5 (c) reducing an intermediate (VII-a) or (VII-b) to obtain an intermediate (VIII-a) or (VIII-b) and subsequently oxidizing the alcohol group in (VIII-a) or (VIII-b) with a mild oxidant to obtain an intermediate (IX-a) or (IX-b) and subsequently alkylating (IX-a) or (IX-b) to obtain (I-f-1) or (I-f-2), which is further alkylated to obtain (I-g-1) or (I-g-2) as in the following reaction schemes:


10




-90-




- 5 (d) converting the alcohol group in (VIII-a) or (VIII-b) to a leaving group and subsequently reacting the thus obtained products with an amine thus obtaining (I-g-1) or (I-g-2)





-91-




- (e) converting an intermediate (IX-a) or (IX-b) to a compound (I-g-1) or (I-g-2) using a Wittig or Wittig-Horner procedure; selectively reducing the double bond in (I-g-1) or (I-g-2) thus obtaining compounds (I-i-1) or (I-i-2); reducing the cyano group in (I-i-1) or (I-i-2) to a methyleneamine group thus obtaining (I-j-1) or (I-j-2); mono- or dialkylating the latter thus obtaining compounds (I-k-1) or (I-k-2); (I-l-1) or (I-l-2):




10



-92-



-93-



and optionally converting the thus obtained compounds of formula (I) into their pharmaceutically acceptable base-addition or acid addition salt form by treatment with a suitable base or acid and conversely treating the base-addition or acid addition salt form with an acid or a base to obtain the free form of the compound of formula (I).

34. A compound of formula (VII-a), (VII-b), (VIII-a), (VIII-b), (IX-a), (IX-b), (I-f-1), (I-f-2), (I-g-1) or (I-g-2) said formula being as in claim 33, wherein G, R¹, R^{2a}, R^{2b}, R³, R⁵, R^{8a}, R^{8b}, R¹² are as claimed in claim 1, and wherein Q is pyrrolidinyl, piperidinyl or homopiperidinyl, substituted on their nitrogen with a radical R⁶ which is C₁₋₆alkyl optionally substituted with one or two, substituents each independently selected from the group consisting of trifluoromethyl, C₃₋₇cycloalkyl, Ar², hydroxy, C₁₋₄alkoxy, C₁₋₄alkylthio, Ar²-oxy-, Ar²-thio-, Ar²(CH₂)_noxy, Ar²(CH₂)_nthio, hydroxycarbonyl, aminocarbonyl, C₁₋₄alkylcarbonyl, Ar²carbonyl, C₁₋₄alkoxycarbonyl, Ar²(CH₂)_ncarbonyl, aminocarbonyloxy, C₁₋₄alkylcarbonyloxy, Ar²carbonyloxy, Ar²(CH₂)_ncarbonyloxy, C₁₋₄alkoxycarbonyl(CH₂)_noxy, mono- or di(C₁₋₄alkyl)aminocarbonyl, mono- or di(C₁₋₄alkyl)aminocarbonyloxy, aminosulfonyl, mono- or di(C₁₋₄alkyl)aminosulfonyl or a heterocycle selected from the group consisting of pyrrolidinyl, pyrrolyl, dihydropyrrolyl, imidazolyl, triazolyl, piperidinyl, homopiperidinyl, piperazinyl, dioxolanyl, dioxanyl, pyridyl and tetrahydropyridyl, wherein each of said heterocycle may optionally be substituted with one or two substituents selected from oxo or C₁₋₆alkyl; and wherein said R⁶ can be represented by R^{6b}, as well as the pharmaceutically acceptable salt forms thereof, and the possible stereoisomeric forms thereof.
35. A compound according to claim 34 wherein R^{6b} is C₁₋₆alkyl optionally substituted with Ar², hydroxy, aminocarbonyl, aminosulfonyl, or C₁₋₆alkyl substituted with two hydroxy radicals, or C₁₋₆alkyl substituted with pyrrolidinyl, piperidinyl, piperazinyl, 4-C₁₋₆alkyl-piperazinyl.
36. A compound according to claim 34 wherein R^{6b} is C₁₋₆alkyl.

37. A compound formula (VII-a), (VII-b), (VIII-a), (VIII-b), (IX-a),
(IX-b), (I-f-1), (I-f-2), (I-g-1) or (I-g-2) said formula being as in claim 33,
wherein G, R¹, R^{2a}, R^{2b}, R³, R⁵, R^{8a}, R^{8b} and R¹² are as claimed in claim 1 and
5 wherein Q is R^{6b} wherein R^{6b} is as claimed in claim 1.
38. A compound according to claim 37 wherein R^{6b} is C₁₋₆alkyl optionally substituted
with Ar², hydroxy, aminocarbonyl, aminosulfonyl, or C₁₋₆alkyl substituted with
two hydroxy radicals, or C₁₋₆alkyl substituted with pyrrolidinyl, piperidinyl,
10 piperazinyl, 4-C₁₋₆alkyl-piperazinyl.
39. A compound according to claim 37 wherein R^{6b} is C₁₋₆alkyl.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/053618

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C07D401/14 A61K31/4184 A61P11/00 A61P31/12 C07D401/06 C07D405/14 //(C07D401/14,235:00,213:00,211:00)		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07D A61K A61P		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, WPI Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 01/00611 A (JANSSENS FRANS EDUARD ; SOMMEN FRANCOIS MARIA (BE); ANDRIES KOENRAA) 4 January 2001 (2001-01-04) cited in the application	1-33
X	page 49, line 4; tables 5,12 -----	34-39
Y	WO 01/00612 A (JANSSENS FRANS EDUARD ; SOMMEN FRANCOIS MARIA (BE); ANDRIES KOENRAA) 4 January 2001 (2001-01-04) cited in the application the whole document -----	1-39
Y	WO 01/00615 A (JANSSENS FRANS EDUARD ; ANDRIES KOENRAAD JOZEF LODENWI (BE); JANSSE) 4 January 2001 (2001-01-04) cited in the application the whole document -----	1-39
-/--		
<div style="display: flex; justify-content: space-between;"> <input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex. </div>		
<div style="display: flex;"> <div style="flex: 1;"> <p>* Special categories of cited documents :</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="flex: 1;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search <div style="text-align: center; font-weight: bold;">8 April 2005</div>		Date of mailing of the international search report <div style="text-align: center; font-weight: bold;">15/04/2005</div>
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer <div style="text-align: center; font-weight: bold;">Lauro, P</div>

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/053618

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 02/092575 A (TRIMERIS INC) 21 November 2002 (2002-11-21) page 67; table 8 -----	1-39
X	WO 02/090347 A (DAISOW CO LTD ;HINOUE KAZUMASA (JP); INOUE YOSHIKAZU (JP); MIKAMI) 14 November 2002 (2002-11-14) page 118; example 84 -----	1-39
A	WO 03/053939 A (THURIEAU CHRISTOPHE ;POITOUT LYDIE (FR); BRAULT VALERIE (FR); FERR) 3 July 2003 (2003-07-03) page 1; examples -----	1-39

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP2004/053618

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0100611	A	04-01-2001	AT 276244 T	15-10-2004
			AU 5816700 A	31-01-2001
			BG 106287 A	31-10-2002
			BR 0012054 A	19-03-2002
			CA 2376781 A1	04-01-2001
			CN 1358180 A	10-07-2002
			CZ 20014574 A3	15-05-2002
			DE 60013836 D1	21-10-2004
			EA 4939 B1	28-10-2004
			EE 200100692 A	17-02-2003
			WO 0100611 A1	04-01-2001
			EP 1196408 A1	17-04-2002
			EP 1418175 A1	12-05-2004
			HR 20010933 A1	30-06-2003
			HU 0201723 A2	28-11-2002
			JP 2003503401 T	28-01-2003
			MX PA02000112 A	02-07-2002
			NO 20016368 A	28-02-2002
			NZ 515418 A	28-11-2003
			PL 362890 A1	02-11-2004
			SI 1196408 T1	28-02-2005
			SK 18942001 A3	08-10-2002
			TR 200103804 T2	21-05-2002
			ZA 200110478 A	20-03-2003
WO 0100612	A	04-01-2001	AT 258928 T	15-02-2004
			AU 778218 B2	25-11-2004
			AU 5816600 A	31-01-2001
			BG 106286 A	31-10-2002
			BR 0012047 A	12-03-2002
			CA 2376676 A1	04-01-2001
			CN 1358181 A	10-07-2002
			DE 60008112 D1	11-03-2004
			DE 60008112 T2	16-09-2004
			DK 1196409 T3	07-06-2004
			EA 5027 B1	28-10-2004
			EE 200100688 A	15-04-2003
			WO 0100612 A2	04-01-2001
			EP 1196409 A2	17-04-2002
			ES 2215683 T3	16-10-2004
			HR 20010935 A1	30-06-2003
			HU 0201869 A2	28-11-2002
			JP 2003503402 T	28-01-2003
			MX PA02000120 A	02-07-2002
			NO 20016369 A	20-02-2002
			NZ 515664 A	30-01-2004
			PL 352378 A1	25-08-2003
			PT 1196409 T	30-06-2004
			SI 1196409 T1	30-06-2004
			SK 19122001 A3	09-01-2003
			TR 200103806 T2	21-06-2002
			US 6747028 B1	08-06-2004
			ZA 200110479 A	20-03-2003
WO 0100615	A	04-01-2001	AT 259796 T	15-03-2004
			AU 774829 B2	08-07-2004
			AU 5222200 A	31-01-2001
			BG 106288 A	31-10-2002

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP2004/053618

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0100615	A		BR 0011997 A CA 2376785 A1 CN 1358182 A ,C CZ 20014573 A3 DE 60008382 D1 DE 60008382 T2 DK 1196410 T3 EA 4746 B1 EE 200100694 A WO 0100615 A1 EP 1196410 A1 EP 1400519 A1 ES 2215670 T3 HR 20010934 A1 HU 0201789 A2 JP 2003503403 T MX PA02000117 A NO 20016370 A NZ 515392 A PL 352385 A1 PT 1196410 T SI 1196410 T1 SK 18952001 A3 TR 200103805 T2 ZA 200110473 A	05-03-2002 04-01-2001 10-07-2002 15-05-2002 25-03-2004 02-12-2004 28-06-2004 26-08-2004 17-02-2003 04-01-2001 17-04-2002 24-03-2004 16-10-2004 30-06-2003 28-11-2002 28-01-2003 02-07-2002 27-12-2001 26-03-2004 25-08-2003 30-07-2004 31-08-2004 06-11-2002 21-06-2002 20-03-2003
WO 02092575	A	21-11-2002	WO 02092575 A1 US 2003119754 A1	21-11-2002 26-06-2003
WO 02090347	A	14-11-2002	EP 1383760 A1 WO 02090347 A1 JP 2004532856 T US 2004157866 A1	28-01-2004 14-11-2002 28-10-2004 12-08-2004
WO 03053939	A	03-07-2003	FR 2833948 A1 AU 2002365016 A1 CA 2471044 A1 CZ 20040819 A3 EP 1467974 A1 WO 03053939 A1 US 2005049290 A1	27-06-2003 09-07-2003 03-07-2003 16-03-2005 20-10-2004 03-07-2003 03-03-2005